

A GRÖBNER BASIS FOR KAZHDAN-LUSZTIG IDEALS

ALEXANDER WOO AND ALEXANDER YONG

ABSTRACT. *Kazhdan–Lusztig ideals*, a family of generalized determinantal ideals investigated in [Woo–Yong ‘08], provide an explicit choice of coordinates and equations encoding a neighbourhood of a torus-fixed point of a Schubert variety on a type A flag variety. Our main result is a Gröbner basis for these ideals. This provides a single geometric setting to transparently explain the naturality of pipe dreams on the *Rothe diagram of a permutation*, and their appearance in:

- combinatorial formulas [Fomin–Kirillov ‘94] for Schubert and Grothendieck polynomials of [Lascoux–Schützenberger ‘82];
- the equivariant K-theory specialization formula of [Buch–Rimányi ‘04]; and
- a positive combinatorial formula for multiplicities of Schubert varieties in good cases, including those for which the associated Kazhdan–Lusztig ideal is homogeneous under the standard grading.

Our results generalize (with alternate proofs) [Knutson–Miller ‘05]’s Gröbner basis theorem for Schubert determinantal ideals and their geometric interpretation of the monomial positivity of Schubert polynomials. We also complement recent work of [Knutson ‘08 & ‘09] on degenerations of Kazhdan–Lusztig varieties in general Lie type, as well as work of [Goldin ‘01] on equivariant localization and of [Lakshmibai–Weyman ‘90], [Rosenthal–Zelevinsky ‘01], and [Krattenthaler ‘01] on Grassmannian multiplicity formulas.

CONTENTS

1. Introduction	2
1.1. Organization of the paper	3
2. Kazhdan–Lusztig ideals and the main theorem	4
2.1. Flag, Schubert and Kazhdan–Lusztig varieties	4
2.2. A choice of coordinate system, equations, and Kazhdan–Lusztig ideals	5
2.3. A Gröbner basis for Kazhdan–Lusztig ideals	6
3. Pipe dreams on Rothe diagrams and the initial ideal of $I_{v,w}$	9
3.1. The prime decomposition theorem for $I_{v,w}$	9
3.2. Pipe complexes are subword complexes	12
3.3. The interlacing strands of pipe dreams	13
4. (Un)specializing Grothendieck polynomials	15
4.1. Four cases of torus actions and their weights	15
4.2. Various graded K-polynomials and multidegrees	16

Date: September 2, 2009.

2000 Mathematics Subject Classification. 14M15, 14N15.

Key words and phrases. Schubert varieties, Gröbner basis, determinantal ideals, equivariant K-theory localization.

4.3. Unspecialized Grothendieck polynomials as K-polynomials	17
4.4. A rationale for unspecializing	22
5. Multiplicities of Schubert varieties	22
5.1. Homogeneity and parabolic moving	23
5.2. Computational results and Monte Carlo simulation	24
5.3. Formulas in the Grassmannian case	26
6. Proof of Theorems 2.1 and 3.2	30
6.1. Vertex decompositions of simplicial complexes	31
6.2. The Kostant–Kumar recursion and the proof of Theorem 6.2	34
6.3. Proof of Theorem 6.1	36
Acknowledgements	40
References	40

1. INTRODUCTION

In our study [WooYon08] of singularities of Schubert varieties, we investigated *Kazhdan–Lusztig ideals* as an explicit choice of coordinates and equations for a neighbourhood of a type A Schubert variety at a torus fixed point. The following problem naturally arises:

Problem 1.1. *Determine a Gröbner basis for the Kazhdan–Lusztig ideals.*

The main result of this paper solves Problem 1.1 by giving an explicit Gröbner basis with squarefree lead terms, thereby deriving a single degeneration from the (reduced and irreducible) variety of the Kazhdan–Lusztig ideal to a reduced union of coordinate subspaces. The class of Kazhdan–Lusztig ideals includes determinantal ideals and many of their generalizations. Gröbner bases for determinantal ideals and their generalizations have been of interest, as their study requires and has applications involving simultaneously commutative algebra, algebraic geometry, representation theory, and combinatorics. Various aspects of determinantal ideals and their generalizations have been extensively treated in the books [BruVet88, MilStu05, Mir07].

We use our Gröbner degeneration to give new explanations for some Schubert combinatorics. In particular, we point out a geometric origin for various combinatorial formulas for Schubert and Grothendieck polynomials [LasSch82a, LasSch82b]. Specifically, we explain the combinatorics of *pipe dreams* on the *Rothe diagram* of a permutation, rather than just the $n \times n$ grid itself, which was the focus of earlier work [KnuMil05, KnuMilYon05]. We thus obtain a geometric explanation and new proof of the formula of A. Buch and R. Rimányi [BucRim04] for the equivariant K-theory relevant specializations of these polynomials (as well as for older formulas of [FomKir94]). This answers a question raised in their paper.

Moreover, our results generalize both the earlier Gröbner geometry explanation of the monomial positivity of Schubert polynomials from [KnuMil05, Theorem A] as well as their companion Gröbner basis theorem [KnuMil05, Theorem B] for *Schubert determinantal ideals*. In brief, as suggested in [Ful92], Schubert determinantal ideals can be realized

as special cases of Kazhdan–Lusztig ideals, allowing us to use recursions for equivariant K-theory Schubert classes due to B. Kostant and S. Kumar [KosKum90]. Although this approach assumes older background, it conceptually streamlines proofs by allowing induction among a larger class of ideals.

This paper is related to developments in [KnuMil04], [KnuMilYon05], [Knu08], and [Knu09]. Specifically, our results complement recent work of A. Knutson [Knu08] on reduced degenerations of *Kazhdan–Lusztig varieties* [KazLus79] for generalized flag varieties of arbitrary Lie type. His paper provided inspiration for the present one. Employing the *limit branchvariety* methods of [Knu07] and *geometric vertex decompositions* [KnuMilYon05], he gives many options for iteratively degenerating a Kazhdan–Lusztig variety into a reduced union of affine spaces; this union is described by a *subword complex* [KnuMil04]. More recently, Knutson also describes in [Knu09] an alternate approach to this degeneration using Frobenius splitting and Bott–Samelson coordinates, though he does not explicitly provide equations.

By comparison and contrast, we work only in type A and single out a specific choice of coordinates and a specific Gröbner degeneration, which we view as especially natural in light of the aforementioned Schubert polynomial combinatorics. Our Gröbner basis theorem provides an explicit, non-iterative realization of one of the degenerations of [Knu08] and [Knu09], using explicit coordinates and explicit equations in those coordinates. However, our techniques are distinct from those of Knutson’s aforementioned work, though parallel to them. We do not use [Knu07], [Knu09], nor specifically any of the new results from [Knu08], substituting instead direct combinatorial commutative algebra arguments. Moreover, our Gröbner basis theorem seems difficult to explicitly adapt in general to other degenerations in [Knu08] and [Knu09], highlighting that the choice of coordinates and term order is somewhat delicate. That the combinatorics lines up with an appropriate choice indicates the resonance between the geometry and the combinatorics.

We expect our methods to extend to finding a generalization of our main result to other Lie types; we hope to address this in a future paper. However, type A warrants some special attention since we would like to further our understanding of the geometric genesis of the beautiful properties of Schubert polynomial and pipe dream combinatorics. As noted in [FomKir96, Section 10], the ensemble of their many properties is specific to type A.

1.1. Organization of the paper. This paper has three central, closely interconnected results (Theorems 2.1, 3.2 and 4.5) and an application, as summarized below.

In Section 2, we recall preliminaries about Kazhdan–Lusztig ideals and state the main result (Theorem 2.1), our Gröbner basis theorem.

In Section 3, we recall *pipe dreams on Rothe diagrams* [FomKir94, BucRim04]. Our second main result, Theorem 3.2, is that they transparently index components of the prime decomposition of the initial ideal (under our term order) of a Kazhdan–Lusztig ideal. We present the associated Reisner–Stanley simplicial complex with faces labeled by these pipe dreams. This complex has the structure of an abstract subword complex [KnuMil04] and thus inherits good properties we use; in particular, it is homeomorphic to a vertex decomposable (and hence shellable) ball or sphere. That said, an additional advantage of using pipe dreams is that they admit an even more “graphical” description as interlacing strand diagrams, akin to the original objects of [FomKir94]; this is not seen at the abstract

subword level. The proof of Theorem 2.1 requires the definitions but not the results of this section.

In Section 4, we introduce *unspecialized Grothendieck polynomials*, a geometrically motivated combinatorial synthesis of results of [FomKir94, BucRim04, KnuMil05]. In terms of these polynomials, our third main result (Theorem 4.5) shows that their specializations to Schubert and Grothendieck polynomials and thus the formulas of [FomKir94, BucRim04] arise from the Gröbner geometry and combinatorial commutative algebra as multidegrees and K-polynomials as defined in, for example, [KnuMil05, MilStu05]. We prove this result assuming Theorem 2.1. Part of our work in this section is to give a proof of a theorem previously known as folklore giving a geometric interpretation of the aforementioned specializations in terms of equivariant K-theory on the flag variety. As with Section 3, the proof of Theorem 2.1 requires some definitions but not the results from this section.

In Section 5, we return to our initial motivation in [WooYon08] of understanding invariants of singularities of Schubert varieties. We relate our work to the open problem of finding an explicit, nonrecursive combinatorial rule for the multiplicity of a Schubert variety at a torus fixed point. We show that in good cases, including those when the Kazhdan–Lusztig ideal is homogeneous under the standard grading, one can positively calculate the multiplicity by counting the pipe dreams of Section 3. In particular, this leads to a simple proof of a determinantal formula, related to but different from previously known formulas [LakWey90, RosZel01], for multiplicities on the Schubert varieties on Grassmannians. Beyond the Grassmannian case, counting pipe dreams solves 98.5% of all multiplicity problems for $n = 5$. Moreover, we point out the efficacy of using of random sampling methods in our study of Schubert varieties, complementing the computational commutative algebra methods of [WooYon08]. Monte Carlo simulation suggests that these cases cover approximately 94% of all multiplicity problems for $n = 6$ and still close to half of all such problems for $n = 10$. We also use this technique to endorse a related, new conjecture.

Finally in Section 6 we give proofs for Theorems 2.1 and 3.2.

2. KAZHDAN–LUSZTIG IDEALS AND THE MAIN THEOREM

2.1. Flag, Schubert and Kazhdan–Lusztig varieties. Let $G = \mathrm{GL}_n(\mathbb{C})$, B be the Borel subgroup of strictly upper triangular matrices, $T \subset B$ the maximal torus of diagonal matrices, and B_- the corresponding opposite Borel subgroup of strictly lower triangular matrices. The **complete flag variety** is

$$\mathrm{Flags}(\mathbb{C}^n) := G/B.$$

The fixed points of G/B under the left action of T are naturally indexed by the symmetric group S_n in its role as the Weyl group of G ; we denote these points e_v for $v \in S_n$. One has a cell decomposition

$$G/B = \coprod_{w \in S_n} B e_w B / B$$

known as the **Bruhat decomposition**. The B -orbit

$$X_w^\circ := B e_w B / B$$

is a cell known as the **Schubert cell**, and its closure

$$X_w := \overline{X_w^\circ}$$

is the **Schubert variety**. It is a subvariety of dimension $\ell(w)$, where $\ell(w)$ is the length of any reduced word of w . Each Schubert variety X_w is a union of Schubert cells; **Bruhat order** is the partial order on S_n defined by declaring that

$$v \leq w \text{ if } X_v^\circ \subseteq X_w.$$

Since every point on X_w is in the B -orbit of some e_v (for $v \leq w$ in Bruhat order), it follows that the study of local questions on Schubert varieties reduces to the case of these fixed points. An affine neighbourhood of e_v is given by $v\Omega_{\text{id}}^\circ$, where in general

$$\Omega_u^\circ := B_- u B / B$$

is known as an **opposite Schubert cell**. Hence to study X_w locally at e_v one only needs to understand $X_w \cap v\Omega_{\text{id}}^\circ$. However, by [KazLus79, Lemma A.4], one has the isomorphism

$$(2.1) \quad X_w \cap v\Omega_{\text{id}}^\circ \cong (X_w \cap \Omega_v^\circ) \times \mathbb{A}^{\ell(v)}.$$

Hence, it is essentially equivalent to study the (reduced and irreducible) **Kazhdan–Lusztig variety**

$$\mathcal{N}_{v,w} = X_w \cap \Omega_v^\circ,$$

forgetting the factor of affine space.

2.2. A choice of coordinate system, equations, and Kazhdan–Lusztig ideals. We now define coordinates on Ω_v° , the **Kazhdan–Lusztig ideal** $I_{v,w}$ in these coordinates, and summarize why this ideal vanishes on $\mathcal{N}_{v,w}$.

Let M_n be the set of all $n \times n$ matrices with entries in $\mathbb{C}$, with coordinate ring $\mathbb{C}[z]$ where $\mathbf{z} = \{z_{i,j}\}_{i,j=1}^n$ are the coordinate functions on the entries of a generic matrix Z . Consistent with our notation in [WooYon08], we index so that $z_{i,j}$ is in the i -th row from the *bottom* of the matrix and j -th column from the left. If we concretely realize G , B , B_- , and T as invertible, upper triangular, lower triangular, and diagonal matrices respectively, then, as explained in [Ful97], we can realize the opposite Schubert cell Ω_v° as an affine subspace of M_n . Specifically, a matrix is in (our realization of) Ω_v° if, for all i ,

$$z_{n-v(i)+1,i} = 1$$

and, for all i ,

$$z_{n-v(i)+1,a} = 0 \text{ and } z_{b,i} = 0 \text{ for } a > i \text{ and } b > n - v(i) + 1.$$

Let

$$\mathbf{z}^{(v)} \subseteq \mathbf{z}$$

denote the remaining unspecialized variables, and $Z^{(v)}$ the specialized generic matrix representing a generic element of Ω_v° .

Example 2.1. If $n = 6$ and $v = 261345$ we have:

$$Z = \begin{pmatrix} z_{61} & z_{62} & z_{63} & z_{64} & z_{65} & z_{66} \\ z_{51} & z_{52} & z_{53} & z_{54} & z_{55} & z_{56} \\ z_{41} & z_{42} & z_{43} & z_{44} & z_{45} & z_{46} \\ z_{31} & z_{32} & z_{33} & z_{34} & z_{35} & z_{36} \\ z_{21} & z_{22} & z_{23} & z_{24} & z_{25} & z_{26} \\ z_{11} & z_{12} & z_{13} & z_{14} & z_{15} & z_{16} \end{pmatrix}, \text{ and } Z^{(v)} = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{41} & 0 & z_{43} & 1 & 0 & 0 \\ z_{31} & 0 & z_{33} & z_{34} & 1 & 0 \\ z_{21} & 0 & z_{23} & z_{24} & z_{25} & 1 \\ z_{11} & 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

□

The ideal $I_{v,w}$ will be an ideal in the polynomial ring $\mathbb{C}[z^{(v)}]$. To describe $I_{v,w}$, let $Z_{ab}^{(v)}$ denote the southwest $a \times b$ submatrix of $Z^{(v)}$. We also let

$$R^w = [r_{ij}^w]_{i,j=1}^n$$

be the **rank matrix** (which we index similarly) defined by

$$r_{ij}^w = \#\{k \mid w(k) \geq n - i + 1, k \leq j\}.$$

Example 2.2. If $w = 365124 \in S_6$ then

$$w = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } R^w = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 3 & 4 & 5 \\ 1 & 2 & 3 & 3 & 3 & 4 \\ 0 & 1 & 2 & 2 & 2 & 3 \\ 0 & 1 & 2 & 2 & 2 & 2 \\ 0 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}.$$

□

Define the **Kazhdan–Lusztig ideal** $I_{v,w}$ to be the ideal of $\mathbb{C}[z^{(v)}]$ generated by all of the size $1 + r_{ij}^w$ minors of $Z_{ij}^{(v)}$ for all i and j . As explained by W. Fulton [Ful92], these minors defining $I_{v,w}$ must vanish on any point of the Schubert variety X_w (no matter what coset representative is chosen to write the point). Therefore, $I_{v,w}$ vanishes on $\mathcal{N}_{v,w}$. We explained in [WooYon08] that $\mathcal{N}_{v,w}$ is in fact defined scheme-theoretically by $I_{v,w}$. Our argument there required the result of Fulton that these minors actually suffice to define (scheme-theoretically) the matrix Schubert variety. However, using only the theorem that $I_{v,w}$ vanishes on $\mathcal{N}_{v,w}$, the arguments of this paper give an independent proof of scheme-theoretic equality, by establishing an equality of Hilbert series (Theorem 6.2).

Not all these minors are needed to generate $I_{v,w}$. The well-known **essential set** for w , denoted $\mathcal{E}(w)$, is recalled in Section 3.1; see (3.1) and (3.2). The **essential minors** are the size $1 + r_{ij}^w$ minors of $Z_{ij}^{(v)}$ for $(i, j) \in \mathcal{E}(w)$. It follows from [Ful92] that this smaller subset of the defining minors of $I_{v,w}$ also generates this ideal.

2.3. A Gröbner basis for Kazhdan–Lusztig ideals. Let $\prec$ be the pure lexicographic term order on monomials in $z^{(v)}$ induced by favoring variables further to the right in $Z^{(v)}$, breaking ties by favoring variables further down in a given column. In symbols,

$$z_{ij} \prec z_{kl} \text{ if } j < l, \text{ or if } j = l \text{ and } i < k.$$

(We use the convention that the leading term is the one which is largest in our order.) For instance, in Example 2.1 we have

$$z_{25} \succ z_{24} \succ z_{34} \succ z_{23} \succ z_{33} \succ z_{43} \succ z_{11} \succ z_{21} \succ z_{31} \succ z_{41}.$$

We now state our main result, which holds for a field $\mathbb{k}$ of arbitrary characteristic, although our proof exploits theorems proved only over $\mathbb{C}$.

Main Theorem 2.1. *Under the term order $\prec$, the essential (and therefore defining) minors form a Gröbner basis for $I_{v,w} \subseteq \mathbb{k}[\mathbf{z}^{(v)}]$.*

Example 2.3. Continuing Examples 2.1 and 2.2, the set of essential minors with respect to Z consists of three 1×1 minors, ten 3×3 minors and five 4×4 minors, corresponding to the three essential set boxes associated to w . They are:

$$\begin{aligned} & z_{11}, z_{21}, z_{31}, \\ & \begin{vmatrix} z_{31} & z_{32} & z_{33} \\ z_{21} & z_{22} & z_{23} \\ z_{11} & z_{12} & z_{13} \end{vmatrix}, \begin{vmatrix} z_{31} & z_{32} & z_{34} \\ z_{21} & z_{22} & z_{24} \\ z_{11} & z_{12} & z_{14} \end{vmatrix}, \dots, \begin{vmatrix} z_{33} & z_{34} & z_{35} \\ z_{23} & z_{24} & z_{25} \\ z_{13} & z_{14} & z_{15} \end{vmatrix}, \text{ and} \\ & \begin{vmatrix} z_{41} & z_{42} & z_{43} & z_{44} \\ z_{31} & z_{32} & z_{33} & z_{34} \\ z_{21} & z_{22} & z_{23} & z_{24} \\ z_{11} & z_{12} & z_{13} & z_{14} \end{vmatrix}, \begin{vmatrix} z_{51} & z_{52} & z_{53} & z_{54} \\ z_{31} & z_{32} & z_{33} & z_{34} \\ z_{21} & z_{22} & z_{23} & z_{24} \\ z_{11} & z_{12} & z_{13} & z_{14} \end{vmatrix}, \dots, \begin{vmatrix} z_{51} & z_{52} & z_{53} & z_{54} \\ z_{41} & z_{42} & z_{43} & z_{44} \\ z_{31} & z_{32} & z_{33} & z_{34} \\ z_{21} & z_{22} & z_{23} & z_{24} \end{vmatrix}. \end{aligned}$$

If we consider the essential minors with respect to $Z^{(261345)}$ we have

$$\begin{aligned} & z_{11}, z_{21}, z_{31}, \\ & \begin{vmatrix} z_{31} & 0 & z_{33} \\ z_{21} & 0 & z_{23} \\ z_{11} & 1 & 0 \end{vmatrix} = -z_{23}z_{31} + z_{33}z_{21}, \begin{vmatrix} z_{31} & 0 & z_{34} \\ z_{21} & 0 & z_{24} \\ z_{11} & 1 & 0 \end{vmatrix}, \dots, \begin{vmatrix} z_{33} & z_{34} & 1 \\ z_{23} & z_{24} & z_{25} \\ 0 & 0 & 0 \end{vmatrix} = 0, \text{ and} \\ & \begin{vmatrix} z_{41} & 0 & z_{43} & 1 \\ z_{31} & 0 & z_{33} & z_{34} \\ z_{21} & 0 & z_{23} & z_{24} \\ z_{11} & 1 & 0 & 0 \end{vmatrix} = z_{41}z_{33}z_{24} - z_{41}z_{23}z_{34} - z_{43}z_{31}z_{24} + z_{43}z_{34}z_{21} + z_{31}z_{23} - z_{23}z_{21}, \\ & \begin{vmatrix} 1 & 0 & 0 & 0 \\ z_{31} & 0 & z_{33} & z_{34} \\ z_{21} & 0 & z_{23} & z_{24} \\ z_{11} & 1 & 0 & 0 \end{vmatrix}, \dots, \begin{vmatrix} 1 & 0 & 0 & 0 \\ z_{41} & 0 & z_{43} & 1 \\ z_{31} & 0 & z_{33} & z_{34} \\ z_{21} & 0 & z_{23} & z_{24} \end{vmatrix} = 0. \end{aligned}$$

which form a Gröbner basis with respect to the same term order, for the Kazhdan–Lusztig ideal $I_{261345,365124}$. As illustrated above, some of the generators may be inhomogeneous (with respect to the standard grading), or even equal to 0. $\square$

Our main theorem specializes to the Gröbner basis theorem of [KnuMil05, Theorem B] (also valid for a field $\mathbb{k}$ of arbitrary characteristic). The **Schubert determinantal ideal** I_w is generated by all size $1 + r_{ij}^w$ minors of the southwest $i \times j$ submatrix Z_{ij} of Z , for all i, j . A **diagonal term order** is one which selects the diagonal (meaning northwest to southeast) term in any minor of Z . (Owing to the indexing of Schubert varieties we are using, which is upside-down and transposed from that of [KnuMil05], our diagonal term order is the same as their *antidiagonal* term order. Moreover, our z_{ij} is their z_{ji} .)

Corollary 2.4. *The essential minors of the Schubert determinantal ideal I_w are a Gröbner basis with respect to any diagonal term order.*

The **matrix Schubert variety** $\bar{X}_w$ is the (reduced and irreducible) variety in M_n defined by I_w . Matrix Schubert varieties were introduced in [Ful92]. The proof of the corollary introduces a construction that shows matrix Schubert varieties are special cases of Kazhdan–Lusztig varieties, as noted in the introduction. We will refer again to this construction in Section 4.3.

Proof of Corollary 2.4: Let

$$w_0 \star w_0 \in S_{2n}$$

be the permutation such that

$$(w_0 \star w_0)(i) = w_0^{(n)}(i) \text{ and } (w_0 \star w_0)(i+n) = w_0^{(n)}(i) + n \text{ for } 1 \leq i \leq n$$

and where $w_0^{(n)}$ is the permutation of longest length in S_n . Furthermore, given $w \in S_n$, let

$$w \times 1_n \in S_{2n}$$

be the standard embedding into S_{2n} , where

$$(w \times 1_n)(i) = w(i) \text{ and } (w \times 1_n)(i+n) = i+n \text{ for } 1 \leq i \leq n.$$

Finally, set

$$\hat{w} = w_0^{(2n)}(w \times 1_n) \in S_{2n}.$$

Notice that $Z^{(w_0 \star w_0)}$ only involves the variables z_{ij} for $1 \leq i, j \leq n$. Now observe that $I_{w_0 \star w_0, \hat{w}}$ has the exact same minors as I_w . Equivalently,

$$\bar{X}_w \cong \mathcal{N}_{w_0 \star w_0, \hat{w}}.$$

Under the term order $\prec$, the diagonal term of any minor is the leading term as long as it is nonzero. Hence by Theorem 2.1, the diagonal terms $\{d_1, \dots, d_N\}$ of the essential minors of $I_{w_0 \star w_0, \hat{w}}$ generate the initial ideal of $I_{w_0 \star w_0, \hat{w}}$ under $\prec$, denoted $\text{in}_{\prec} I_{w_0 \star w_0, \hat{w}}$. For the term order $\prec'$, the result now follows as a special case of Theorem 2.1.

Now suppose $\prec'$ is another diagonal term order. Then if $\text{in}_{\prec'} I_{w_0 \star w_0, \hat{w}}$ is the initial ideal with respect to $\prec'$, we have

$$\text{in}_{\prec} I_{w_0 \star w_0, \hat{w}} = \langle d_1, \dots, d_N \rangle \subseteq \text{in}_{\prec'} I_{w_0 \star w_0, \hat{w}}.$$

However, since both $\text{in}_{\prec} I_{w_0 \star w_0, \hat{w}}$ and $\text{in}_{\prec'} I_{w_0 \star w_0, \hat{w}}$ are Gröbner degenerations of $I_{w_0 \star w_0, \hat{w}}$, they have the same Hilbert series (under the appropriate grading as described in Section 4.1). Hence they are equal and the theorem holds for $\prec'$. $\square$

Example 2.5. Let $w = 2143 \in S_4$. Then since $n = 4$, we have

$$w_0 \star w_0 = 43218765 \in S_8, \quad w \times 1_4 = 21435678, \text{ and } \hat{w} = w_0^{(8)}(w \times 1_4) = 78564321.$$

Hence

$$Z^{(w_0 \star w_0)} = Z^{(43218765)} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ z_{41} & z_{42} & z_{43} & z_{44} & 0 & 0 & 0 & 1 \\ z_{31} & z_{32} & z_{33} & z_{34} & 0 & 0 & 1 & 0 \\ z_{21} & z_{22} & z_{23} & z_{24} & 0 & 1 & 0 & 0 \\ z_{11} & z_{12} & z_{13} & z_{14} & 1 & 0 & 0 & 0 \end{pmatrix}.$$

The reader can check that

$$I_{w_0 \star w_0, \hat{w}} = \left\langle z_{11}, \begin{vmatrix} z_{31} & z_{32} & z_{33} \\ z_{21} & z_{22} & z_{23} \\ z_{11} & z_{12} & z_{13} \end{vmatrix} \right\rangle,$$

which are the essential minors of I_{2143} . (Compare with the first example of [KnuMil05].) $\square$

3. PIPE DREAMS ON ROTHE DIAGRAMS AND THE INITIAL IDEAL OF $I_{v,w}$

3.1. The prime decomposition theorem for $I_{v,w}$. Let $\text{LT}_{\prec}(f)$ denote the leading term of f under the order $\prec$. The **initial ideal**

$$\text{in}_{\prec} I_{v,w} = \langle \text{LT}_{\prec}(f) : f \in I_{v,w} \rangle$$

is by virtue of Theorem 2.1 a squarefree monomial ideal. To help understand this initial ideal, we use the **Reisner–Stanley correspondence**, a bijective correspondence between squarefree monomial ideals and simplicial complexes.

Recall that given a simplicial complex Δ on a vertex set A (so our faces are subsets of A), the **Reisner–Stanley ideal** $I(\Delta)$ is the squarefree monomial ideal in

$$S = \mathbb{k}[x_a \mid a \in A]$$

generated by monomials corresponding to faces not in Δ . In symbols,

$$I(\Delta) = \left\langle \prod_{a \in F} x_a \mid F \notin \Delta \right\rangle.$$

The **Reisner–Stanley ring** of Δ is defined to be $S/I(\Delta)$. Conversely, given a squarefree monomial ideal I in the ring S , we can associate to it the **Reisner–Stanley complex** $\Delta(I)$, which is the simplicial complex on A such that

$$F \in \Delta(I) \text{ if } \prod_{a \in F} x_a \notin I.$$

We now give a combinatorial description of a simplicial complex $\Delta_{v,w}$ which we call the **pipe complex**. We will show that $\Delta_{v,w}$ is the Reisner–Stanley complex of $\text{in}_{\prec} I_{v,w}$.

We give coordinates to an ambient $n \times n$ grid so that $(1, 1)$ refers to the southwest corner, $(n, 1)$ refers to the northwest corner, and so on. Now, to each $v \in S_n$, the **Rothe diagram** $D(v)$ is the following subset of the $n \times n$ grid:

$$(3.1) \quad D(v) = \{(i, j) : i < n - w(j) + 1 \text{ and } j < w^{-1}(n - i + 1)\}.$$

Alternatively, this set is described as follows. Place a dot $\bullet$ in position $(n - w(j) + 1, j)$ for $1 \leq j \leq n$. For each dot draw the “hook” that extends to the right and above that dot. The boxes that are not in any hook are the boxes of $D(v)$.

Notice that with these conventions, the coordinates of the boxes of $D(v)$ are exactly the labels for the (unspecialized) variables appearing in $Z^{(v)}$.

The **essential set** $\mathcal{E}(v)$ can be described as the set of those boxes which are on the north-east edge of some connected components of $D(v)$. To be precise,

$$(3.2) \quad (i, j) \in \mathcal{E}(v) \text{ if } (i, j) \in D(v) \text{ but both } (i + 1, j) \notin D(v) \text{ and } (i, j + 1) \notin D(v).$$

The **canonical labeling** of $D(v)$ is obtained by filling the t boxes in row i with the labels $i, i+1, \dots, i+t-1$ (with i being the label of the leftmost box). We will interchangeably refer to a box in $D(v)$ with label ℓ and a generator u_ℓ of the **NilHecke algebra** $\mathcal{A}_n$. The NilHecke algebra is the $\mathbb{C}$ -algebra with generators $u_1, \dots, u_{n-1}$ and relations:

$$(3.3) \quad \begin{aligned} u_i u_j &= u_j u_i, \quad |i-j| \geq 2; \\ u_i u_{i+1} u_i &= u_{i+1} u_i u_{i+1}; \\ u_i^2 &= u_i. \end{aligned}$$

(The usual presentation has the relation $u_i^2 = -u_i$, but this equivalent presentation simplifies the sequel.) The NilHecke algebra $\mathcal{A}_n$ has a basis $\{u_w\}$ indexed by permutations $w \in S_n$. For the simple transposition

$$s_i := (i \leftrightarrow i+1),$$

let

$$u_{s_i} := u_i,$$

and if

$$w = s_{i_1} \cdots s_{i_\ell}$$

is any reduced transposition for w , then let

$$u_w := u_{i_1} \cdots u_{i_\ell}.$$

Multiplication in this basis by a generator u_i is as follows.

$$u_w u_i = \begin{cases} u_w & \text{if } ws_i < w; \\ u_{ws_i} & \text{otherwise.} \end{cases}$$

A **pipe dream** on $D(v)$ is a configuration $\mathcal{P}$ of crosses $+$ in a subset of boxes of $D(v)$. Let $\text{Pipes}(v)$ be the set of all such pipe dreams. Associated to $\mathcal{P}$ is the **Demazure product** $\prod \mathcal{P}$ of generators u_ℓ obtained by reading the crosses in rows, from left to right, and then from top to bottom. Fix $w \in S_n$. Following [BucRim04], a pipe dream $\mathcal{P}$ on $D(v)$ is furthermore a **pipe dream for w** if

$$\prod \mathcal{P} = u_w$$

Such a pipe dream is **reduced** if

$$\#\mathcal{P} = \ell(w).$$

Let

$$\text{Pipes}(v, w) \subseteq \text{Pipes}(v)$$

be the collection of all pipe dreams on $D(v)$ for w , and let


$$\text{RedPipes}(v, w) \subseteq \text{Pipes}(v, w)$$

be the subset of reduced ones. Finally, note that if $\mathcal{P}$ is the pipe dream with a $+$ in every box of $D(v)$ then

$$\prod \mathcal{P} = u_{w_0 v}.$$

This reformulates the well known fact that the canonical labeling of $D(v)$ encodes, via the reading order we use, a reduced word of $w_0 v$; see for example [Man01, Remark 2.1.9].

Example 3.1. Let $v = 31524$ (in one line notation). Then the diagram $D(v)$ and the associated canonical labeling is given below:




Here,


$$\mathcal{E}(v) = \{(2, 2), (4, 2), (2, 4)\}.$$

The reader can check that the canonical labeling gives a reduced word $s_4 s_2 s_3 s_4 s_1 s_2$ for $w_0 v$.

Let $w = 13254 = s_2 s_4 = s_4 s_2$. Then $\text{RedPipes}(v, w)$ consists of:



whereas $\text{Pipes}(v, w)$ consists of the above pipe dreams together with:



Lastly, $\text{Pipes}(v)$ consists of the above pipe dreams together with all remaining $2^6 - 9 = 23$ pipe dreams whose Demazure product does not give w . $\square$

We are now ready to define the pipe complex $\Delta_{v,w}$. The complex $\Delta_{v,w}$ is a complex on the vertex set $D(v)$ such that F is a face if $D(v) \setminus F$ contains a (reduced) pipe dream for $w_0 w$ (not w). Following [KnuMil04] it is convenient for us to label the faces of $\Delta_{v,w}$ by pipe dreams, so that a face F will be labeled by the pipe dream $\mathcal{P}$ with crosses everywhere except at the vertices of F . This means that vertices are labeled by pipe dreams with precisely $\ell(w_0 v) - 1$ $+$'s, and the empty face is labeled by the pipe dream with a $+$ in every square of $D(v)$.

Pipe dreams on $D(v)$ were introduced in [BucRim04] on purely combinatorial grounds. Our main result about $\Delta_{v,w}$ is the following, which gives a geometric rationale for these pipe dreams. Moreover, we will use this result in the next section to geometrically explain the specialization formula of [BucRim04] for Grothendieck polynomials (as well as a generalization).

Theorem 3.2. For $v, w \in S_n$, $\Delta_{v,w}$ is the Reisner–Stanley complex of $\text{in}_{\prec} I_{v,w}$. Therefore, the following prime decomposition holds:

$$(3.4) \quad \text{in}_{\prec} I_{v,w} = \bigcap_{\mathcal{P} \in \text{RedPipes}(v, w_0 w)} \langle z_{ij} \mid (i, j) \in \mathcal{P} \rangle.$$

The proof, basically given in parallel with the proof of Theorem 2.1, is provided in Section 6.

3.2. Pipe complexes are subword complexes. We now state some facts about pipe complexes which will be needed in our proof of Theorem 3.2. To do so, we need to recall some definitions from [KnuMil04]. Let

$$Q = (\alpha_1, \dots, \alpha_\ell)$$

be a sequence from the alphabet $\{1, 2, \dots, n-1\}$ such that if $s_i = (i \leftrightarrow i+1)$ is the simple reflection in S_n then $s_{\alpha_1} \cdots s_{\alpha_\ell}$ is a reduced word for $v \in S_n$. In particular, $\ell = \ell(v)$.

Fix $w \in S_n$ and a word Q in the simple generators $\{s_1, \dots, s_{n-1}\}$. In [KnuMil04], the **subword complex** $\Delta(Q, w)$ is defined to be the simplicial complex where the vertex set is the positions $1, 2, \dots, \ell$ of Q with $F \subseteq \{1, \dots, \ell\}$ defined to be a facet if and only if $\prod_{i \in F} Q \setminus F$, the product of simple roots corresponding to the positions $\{1, \dots, \ell\} \setminus F$, is a reduced word for w . More generally, faces

$$F \in \Delta(Q, w)$$

correspond to subwords of Q such that

$$\prod Q \setminus F \geq w$$


in Bruhat order.

Let us record the following result, which is immediate from results of [KnuMil04] and our definitions.

Proposition 3.3. $\Delta_{v,w}$ is the subword complex $\Delta(Q, w_0 w)$ where Q is the canonical labeling of $D(v)$ in our reading order and therefore a reduced word for $w_0 v$. Hence $\Delta_{v,w}$ is shellable and homeomorphic to a ball or a sphere. Moreover, the facets of $\Delta_{v,w}$ are labeled by $\mathcal{P} \in \text{RedPipes}(v, w_0 w)$ and interior faces labeled by $\mathcal{P} \in \text{Pipes}(v, w_0 w)$.


Proof. From the definition of $\Delta_{v,w}$ and $\Delta(Q, w_0 w)$ the first claim amounts to the well-known fact that the canonical filling encodes a reduced word for $w_0 v$. The remaining results are general properties of any subword complex; see Theorems 2.5 and 3.7 in [KnuMil04]. $\square$

Example 3.4. Let $v = 31542$. Then $D(v)$ and the canonical labeling are given by:



Let $w = 53142$ and hence $w_0 w = 13524 = s_4 s_2 s_3$. Then the pipe complex $\Delta_{v,w} = \Delta_{31542, 53142}$ is the one dimensional ball of Figure 1. $\square$

Example 3.5. Let $v = 31452$ and $w = 53142$. So $w_0 w = 13524 = s_4 s_2 s_3$ as above. We have:






FIGURE 1. $\Delta_{31542,53142}$ is a one-dimensional ball




FIGURE 2. $\Delta_{31452,53142}$ is a two dimensional ball

Therefore, $\Delta_{v,w} = \Delta_{31452,53142}$ is the complex given by Figure 2.

□

3.3. The interlacing strands of pipe dreams. Theorem 3.2 explains the geometric naturality of drawing pipe dreams as we do rather than considering them only abstractly as

subwords of a fixed reduced word. This is closer to the point of view of [KnuMil05] or [KnuMilYon05] than of [KnuMil04] or [Knu08].


Moreover, there is a further combinatorial property of pipe dreams in $\text{Pipes}(v, w)$, also not seen at the subword level, that furthermore justifies the nomenclature in relation to graphical objects of [FomKir94], so named **RC graphs** in [BerBil93] and renamed **pipe dreams** in [KnuMil05]. In those earlier papers, one tiles each square on an $n \times n$ grid by crosses $+$ and elbows $\neg\lrcorner$, resulting in an attractive collection of strands which visibly encode a permutation. Let us explain how we can similarly introduce strands into our pipe dreams.

Given the diagram $D(v)$ of v , let $\text{flatten}(D(v))$ denote the **flattening** of $D(v)$, which is defined by compressing the squares in each column southward past all non-diagram squares. This fixes an obvious bijection between $D(v)$ and $\text{flatten}(D(v))$. Moreover, if $\mathcal{P} \in \text{Pipes}(v, w)$ then we can construct a new pipe dream $\hat{\mathcal{P}}$ by placing a cross $+$ in each square of $\text{flatten}(D(v))$ if and only if a cross $+$ appears in the corresponding square of $D(v)$. Now define $\overline{\mathcal{P}}$ by inserting an elbow $\neg\lrcorner$ in all other 1×1 squares of the $n \times n$ grid.

Proposition 3.6. *If $\mathcal{P} \in \text{RedPipes}(v, w)$, then $\overline{\mathcal{P}}$ consists of pipes such that the strand that starts in column i ends in row $w(i)$ (as counted from the bottom). In addition, no two strands cross more than once, and crosses $+$ occur in the strict lower triangular part.*

Proof. Consider the configuration $\overline{\mathcal{P}}$ on the $n \times n$ grid. Adorn the said grid with the canonical labeling, so $1, 2, \dots, n$ will line the bottom-most row, then $2, 3, \dots, n-1$ will line the next row, etc. Now note that flatten sends each $+$ in $\mathcal{P}$ to a $+$ in $\overline{\mathcal{P}}$ such that the associated labeling under the canonical labelings of $D(v)$ and $n \times n$ is the same. Moreover, the reading words of both pipe dreams is the same. Finally, notice that there are at most $n-i$ boxes of $D(v)$ in column i . Hence $\overline{\mathcal{P}}$ must have all of its $+$'s in the strict lower triangular part of $n \times n$. Therefore, it follows from the discussion found in Section 5 (and specifically of Example 5.1) of [KnuMil04] that $\overline{\mathcal{P}}$ is a pipe dream for w in the sense of [FomKir94], and the proposition follows. $\square$

Example 3.7. Below are the facets $\mathcal{P} \in \text{RedPipes}(31452, 13524)$ of the pipe complex from Figure 2, and their corresponding flattenings $\overline{\mathcal{P}}$.



$\square$

Note that in the special case where $\mathcal{P} \in \text{RedPipes}(w_0 \star w_0, \hat{w})$, flattening $D(w_0 \star w_0)$ does nothing:

$$\text{flatten}(D(w_0 \star w_0)) = D(w_0 \star w_0),$$

and $\hat{\mathcal{P}} = \mathcal{P}$. Then these $\mathcal{P}$ are precisely all the pipe dreams for w (after rotation and reflection to match conventions) in the sense of [FomKir94].

4. (UN)SPECIALIZING GROTHENDIECK POLYNOMIALS

4.1. Four cases of torus actions and their weights. We describe some torus actions on $\text{Fun}[\Omega_v] \cong \mathbb{C}[z^{(v)}]$ we use in this paper.

First and more importantly, the action of $T \cong (\mathbb{C}^*)^n$ on $\text{Flags}(\mathbb{C}^n)$ induces what we will call the **usual action**. This action is the left action of diagonal matrices on B-cosets of G written in our coordinates. The action rescales rows independently and rescales columns dependently, as upon rescaling a row one must rescale a corresponding column to ensure there is a 1 in position $(n - v(j) + 1, j)$ (as read with our usual upside-down matrix coordinates). Adopting the usual convention that the homomorphism picking out the i -th diagonal entry is the weight t_i and writing weights additively, this action gives the matrix entry at (i, j) the weight $t_{n-i+1} - t_{v(j)}$. The variable z_{ij} is the coordinate function on this matrix entry and therefore (the torus action on the variable) has weight

$$\text{wt}(z_{ij}) = t_{v(j)} - t_{n-i+1}.$$

Note that the weight of $Z_{i,j}$ is always a positive root, and hence a positive integer linear combination of the positive simple roots $t_i - t_{i+1}$. This action always fixes the coordinate ring of $\mathcal{N}_{v,w}$.

Second, in the special case of the usual action where

$$v = w_0 \star w_0$$

(as defined after Corollary 2.4), this usual action can be thought of as independently rescaling the bottom n rows and the leftmost n columns, while dependently scaling the remaining rows and columns to fix the matrix outside of the southwest $n \times n$ block. It is convenient in this case to relabel the weights by

$$x_j = t_{n-j+1} = t_{v(j)}, \text{ and } y_i = t_{2n+1-i}.$$

This gives each variable z_{ij} the weight

$$\text{wt}(z_{ij}) = x_j - y_i.$$

In this case, as explained in the proof of Corollary 2.4, the Kazhdan–Lusztig variety $\mathcal{N}_{w_0 \star w_0, \hat{w}}$ is the matrix Schubert variety $\bar{X}_w$ defined in Section 2.3. We will see that the multidegree and K-polynomial of the coordinate ring of $\mathcal{N}_{w_0 \star w_0, \hat{w}}$ with respect to this grading and weight labeling are respectively the double Schubert and double Grothendieck polynomials of [LasSch82a, LasSch82b].

Third, there is the action of $\mathbb{C}^*$ that equally rescales each variable in $\mathbb{C}[z^{(v)}]$ with the same weight t ; this is the **dilation action**. This action fixes $\mathcal{N}_{v,w}$ if $I_{v,w}$ is homogeneous under the standard grading that assigns each variable z_{ij}

$$\text{wt}(z_{ij}) = 1.$$

In this case we say that $I_{v,w}$ is **standardly homogeneous**. As was pointed out to us in a private communication by A. Knutson, this automatically happens if there exists a coweight

λ for which $\langle t_{v(j)} - t_{n-i+1}, \lambda \rangle = 1$ for all i, j where z_{ij} is a variable in $\mathbb{C}[z^{(v)}]$, in which case we say that $w_0 v$ is **λ -cominuscul**. Note that this condition does not depend on w . If we take into account w , there are other cases for which $\mathcal{N}_{v,w}$ is fixed by the dilation action, but we know of no useful characterization; see Section 5.

Fourth and finally, there is the **rescaling action** of $(\mathbb{C}^*)^{\ell(w_0 v)}$ that independently rescales each variable z_{ij} with weight

$$\text{wt}(z_{ij}) = t_{ij}.$$

This action preserves only unions of coordinate subspaces (and other monomial subschemes in our coordinates).

4.2. Various graded K-polynomials and multidegrees. We now use some notions from combinatorial commutative algebra which can be found in the textbook [MilStu05].

Consider a polynomial ring

$$S = \mathbb{C}[z_1, \dots, z_m]$$

with a grading such that z_i has some degree $a_i \in \mathbb{Z}^N$. A finitely graded S -module

$$M = \bigoplus_{v \in \mathbb{Z}^N} M_v$$

over R has a free resolution

$$E_\bullet : 0 \leftarrow E_1 \leftarrow E_2 \leftarrow \dots \leftarrow E_L \leftarrow 0$$

where

$$E_i = \bigoplus_{j=1}^{\beta_j} R(-d_{ij})$$

is graded with the j -th summand of E_i generated in degree $d_{ij} \in \mathbb{Z}^N$.

Then the $(\mathbb{Z}^N$ -graded) **K-polynomial** of M is

$$\mathcal{K}(M, t) = \sum_j (-1)^j \sum_i t^{d_{ij}}.$$

In any case where S is **positively graded**, meaning that the a_i generate a pointed cone in $\mathbb{Z}^N$, $\mathcal{K}(M, t)$ is the numerator of the $\mathbb{Z}^N$ -graded Hilbert series:

$$\text{Hilb}(M, t) = \frac{\mathcal{K}(M, t)}{\prod_i (1 - t^{a_i})}.$$

The **multidegree** $\mathcal{C}(M, t)$ is by definition the sum of the lowest degree terms of $\mathcal{K}(M, 1 - t)$. (This means we substitute $1 - t_k$ for t_k for all k , $1 < k < N$.)

In Section 4.3, the geometric underpinnings of (un)specializing Grothendieck and Schubert polynomials are explained in terms of inclusions of tori. We therefore present a discussion of the necessary background now.

Suppose our grading comes from a group action of $(\mathbb{C}^*)^N$ on S ; this means that the grading group $\mathbb{Z}^N$ is identified with the weight lattice of $(\mathbb{C}^*)^N$, and a_i is the weight (written additively) of the action on z_i . In this case, a quotient ring S/I is a homogeneous S -module under our grading if and only if the affine variety (or scheme) $V(I)$ is fixed by the $(\mathbb{C}^*)^N$ action. Furthermore $\mathcal{K}(S/I, t)$ is now the equivariant K-theory class $[\mathcal{O}_{V(I)}] \in K_T^0(\mathbb{C}^m)$, whereas $\mathcal{C}(R/I, t)$ is the equivariant cohomology class $[V(I)] \in H_T(\mathbb{C}^m)$. Note that, while we wrote our weights additively in describing the degree a_i given to the variable z_i , in

the K-polynomial the weights are “exponentiated” and written as t^{a_i} . Since $\mathbb{C}^m$ is contractible to a point, $K_T^0(\mathbb{C}^m)$ can be identified with the K_T -ring of a point, which is the representation ring of $T = (\mathbb{C}^*)^N$. This ring is isomorphic to the Laurent polynomial ring in k variables, and weights are multiplicative in this ring. On the other hand, essentially due to [MilStu05, Prop. 8.49], weights are additive in the cohomology ring.

Suppose moreover that we have tori

$$T_1 = (\mathbb{C}^*)^M \text{ and } T_2 = (\mathbb{C}^*)^N$$

acting on S , with a map of tori

$$\rho : T_1 \rightarrow T_2$$

which commutes with this action. The map of tori induces a map ρ^* from weights of T_2 to weights of T_1 . This in turn induces a map

$$K_{T_2}^0(\mathbb{C}^m) \rightarrow K_{T_1}^0(\mathbb{C}^m).$$

On the level of K-polynomials, one can obtain the K-polynomial $\mathcal{K}_{T_1}(M, s)$ with respect to the grading from T_1 from the K-polynomial $\mathcal{K}_{T_2}(M, t)$ by substituting $s^{\rho^*(e_i)}$ for each t_i . (Here e_i is the vector with 1 in the i -th coordinate and 0 elsewhere.) On the other hand, weights are additive in cohomology, so one obtains the multidegree $\mathcal{C}_{T_1}(M, s)$ from $\mathcal{C}_{T_2}(M, t)$ by substituting $\langle \rho^*(e_i), s \rangle$ for t_i .

In Section 4.3 below, S is the ring $\mathbb{C}[z^{(v)}]$, considered as the coordinate ring of the affine space $\Omega_v^\circ \cong \mathbb{C}^{\ell(w_0 v)}$, and the torus action will be one of those specified in the previous section. Note that there is an embedding of the (torus for) the usual action into the rescaling action, and, in the λ -cominuscule case, an embedding (by the coweight λ) of the dilation action into the usual action.

4.3. Unspecialized Grothendieck polynomials as K-polynomials. Recall the NilHecke algebra $\mathcal{A}_n$ defined by (3.3) in Section 2. Consider the following generating series

$$\tilde{\mathfrak{G}}_v = \prod_{(i,j) \in D(v)} (1 + u_{\text{label}(i,j)}(1 - t_{ij})) \in \mathcal{A}_n[[t_{ij}]],$$

where $\text{label}(i, j)$ is the label in box (i, j) (in the i -th row from the bottom and j -th column from the left in accordance with our convention), of the canonical labeling of $D(v)$ and where the product is taken from left to right along rows and from top to bottom (in accordance with the reading of the canonical labeling). Similar generating series were considered in [FomKir94, BucRim04].

Now define the **unspecialized Grothendieck polynomial** by

$$\mathfrak{G}_{v,w}(t_{11}, t_{12}, \dots) := \text{coefficient of } w_0 w \text{ in } \tilde{\mathfrak{G}}_v.$$

It is clear from the construction that

$$(4.1) \quad \mathfrak{G}_{v,w}(t_{11}, t_{12}, \dots) = \sum_{\mathcal{P} \in \text{Pipes}(v, w_0 w)} (-1)^{\#\mathcal{P} - \ell(w_0 w)} (1 - t^{\mathcal{P}}),$$

where

$$1 - t^{\mathcal{P}} = \prod_{(i,j) \text{ contains a } +} (1 - t_{ij}).$$

We furthermore define the **unspecialized Schubert polynomial** by

$$(4.2) \quad \mathfrak{S}_{v,w} = \sum_{\mathcal{P} \in \text{RedPipes}(v, w_0 w)} t^{\mathcal{P}}.$$

The following result interprets $\mathfrak{S}_{v,w}(t)$ and $\mathfrak{S}_{v,w}(t)$ in terms of the pipe complex $\Delta_{v,w}$:

Proposition 4.1. *$\mathfrak{S}_{v,w}(t)$ and $\mathfrak{S}_{v,w}(t)$ are respectively the K -polynomial and multidegree of $R/\text{in}_{\prec} I_{v,w}$ under the rescaling action. Equivalently, they are the K -polynomial and multidegree of $R/K_{v,w}$, the Reisner–Stanley ring of $\Delta_{v,w}$. Furthermore, $\mathfrak{S}_{v,w}$ is the lowest degree term of $\mathfrak{S}_{v,w}(1 - t)$. (This is the result of substituting $1 - t_{ij}$ for t_{ij} for all i and j .)*

Proof. From Proposition 3.3, we know $\Delta_{v,w}$ is a ball or sphere. It is established in [KnuMil04, Theorem 4.1] that whenever Δ is a ball or sphere and S is its Reisner–Stanley ring, then

$$\mathcal{K}(S, t) = \sum_F (-1)^{\dim(\Delta) - \dim(F)} \prod_{i \notin F} (1 - t_i),$$

where the sum is over all the *internal* faces F of Δ . (Their statement is for the case of the subword complex although their proof, a short derivation from Hochster’s formula [Hoc77], applies more generally.) Now, $\mathfrak{S}_{v,w}(t)$ is by construction exactly this sum for $\Delta_{v,w}$. (Here our labeling of a face F by the pipe dream with crosses everywhere except the vertices of F makes our statement much cleaner.) This proves our statement for $R/K_{v,w}$. Since the facets of $\Delta_{v,w}$ are precisely the faces labeled by reduced pipe dreams, the multidegree statement for $R/K_{v,w}$ follows [MilStu05, Prop. 8.49, Thm. 8.53].

Theorem 3.2 says $\Delta_{v,w}$ is the Reisner–Stanley complex of $\mathbb{C}[z^{(v)}]/\text{in}_{\prec} I_{v,w}$, so the statements for $R/\text{in}_{\prec} I_{v,w}$ follow. (Alternatively, we can deduce this result directly from Theorem 6.2.)

The stated relationship between $\mathfrak{S}_{v,w}$ and $\mathfrak{S}_{v,w}$ is the standard relationship between K -polynomials and multidegrees; see [MilStu05, Chapter 8.5]. $\square$

Example 4.2. Continuing Example 3.5, we have

$$\begin{aligned} \tilde{\mathfrak{S}}_{31452} = & (1 + u_4(1 - t_{42}))(1 + u_2(1 - t_{21}))(1 + u_3(1 - t_{22}))(1 + u_1(1 - t_{11})) \times \\ & (1 + u_2(1 - t_{12}))(1 + u_3(1 - t_{13})). \end{aligned}$$

Expanding, and collecting all terms with coefficient

$$u_{w_0 w} = u_{13524} = u_4 u_2 u_3 \equiv u_2 u_4 u_3$$

gives an alternating sum over all the internal faces of $\Delta_{31452, 53142}$, namely

$$\begin{aligned} \mathfrak{S}_{31452, 53142}(t) = & (1 - t_{42})(1 - t_{21})(1 - t_{22}) + (1 - t_{42})(1 - t_{21})(1 - t_{13}) + (1 - t_{42})(1 - t_{12})(1 - t_{13}) \\ & - (1 - t_{42})(1 - t_{21})(1 - t_{22})(1 - t_{13}) - (1 - t_{42})(1 - t_{21})(1 - t_{12})(1 - t_{13}). \end{aligned}$$

If we calculate the unspecialized Schubert polynomial, we get

$$\mathfrak{S}_{31452, 53142}(t) = t_{42} t_{21} t_{22} + t_{42} t_{21} t_{13} + t_{42} t_{12} t_{13}.$$

$\square$

Rather than give a standard definition of the double Grothendieck and Schubert polynomials of [LasSch82a, LasSch82b], we prefer from our viewpoint to define them via the unspecialized versions, proving the equivalence by assuming the formula of [FomKir94].

In fact, the final claim of the definition–theorem below recovers [KnuMil05, Theorem A], which states that the double Grothendieck polynomial $\mathfrak{G}_w(\mathbf{x}, \mathbf{y})$ is the K-polynomial of a matrix Schubert variety $\overline{X}_w$ (which is isomorphic to the Kazhdan–Lusztig variety $\mathcal{N}_{w_0 \star w_0, \hat{w}}$ where $w_0 \star w_0$ and $\hat{w}$ are defined as in the proof of Corollary 2.4).

Definition–Theorem 4.3. *The double Grothendieck polynomial and double Schubert polynomial of [LasSch82a, LasSch82b] are given by*

$$(4.3) \quad \mathfrak{G}_w(\mathbf{x}, \mathbf{y}) = \mathfrak{G}_{w_0 \star w_0, \hat{w}}(t_{ij} \mapsto x_j/y_i), \text{ and}$$

$$(4.4) \quad \mathfrak{S}_w(\mathbf{x}, \mathbf{y}) = \mathfrak{S}_{w_0 \star w_0, \hat{w}}(t_{ij} \mapsto x_j - y_i).$$

In particular, these give respectively the K-polynomial and multidegree of $\mathcal{N}_{w_0 \star w_0, \hat{w}}$ under our special case of the usual action.

Example 4.4. Let us compute $\mathfrak{S}_{13524}(\mathbf{x}, \mathbf{y})$. Here $w = 13524$, so $\hat{w} = \underline{10869754321} \in S_{10}$. To compute $\mathfrak{S}_{w_0 \star w_0, \hat{w}}(\mathbf{t})$, we consider all pipe dreams in the 5×5 box given by $D(w_0 \star w_0)$ whose associated product is $w_0^{(10)} \hat{w} = 135246789\underline{10} = w \times 1_5$. The reader can check that there are six such pipe dreams, and summing up their weights under $t_{ij} \mapsto x_i - y_j$ gives, by Definition–Theorem 4.3:

$$\begin{aligned} \mathfrak{S}_{13524}(\mathbf{x}, \mathbf{y}) = & (x_2 - y_3)(x_1 - y_2)(x_2 - y_2) + (x_2 - y_3)(x_1 - y_2)(x_3 - y_1) \\ & + (x_2 - y_3)(x_2 - y_1)(x_3 - y_1) + (x_1 - y_4)(x_1 - y_2)(x_3 - y_1) + (x_1 - y_4)(x_1 - y_2)(x_2 - y_2) \\ & + (x_1 - y_4)(x_2 - y_1)(x_3 - y_1) \end{aligned}$$

□

Proof of Definition–Theorem 4.3: Under our conventions, the Schubert polynomial formula [FomKir94] states

$$\mathfrak{S}_w(\mathbf{x}, \mathbf{y}) = \sum_{\mathcal{P}} \prod_{(i,j) \in \mathcal{P}} x_j - y_i,$$

where the sum is over reduced pipe dreams for w fitting inside an $n \times n$ square. Moreover, in [FomKir94] one also has (under our conventions)

$$\mathfrak{G}_w(\mathbf{x}, \mathbf{y}) = \sum_{\mathcal{P}} (-1)^{\#\mathcal{P} - \ell(w)} \prod_{(i,j) \in \mathcal{P}} 1 - x_j/y_i,$$

with the sum being over all pipe dreams for w on $n \times n$. It is straightforward to check that $\text{Pipes}(w_0 \star w_0, \hat{w})$ is the same set of pipe dreams. (Previous authors write their pipe dreams transposed and turned upside down from ours because they used conventions natural for the study of Schubert polynomials rather than conventions natural for the study of Schubert varieties.) Thus (4.1), after the substitution

$$t_{ij} \mapsto x_j/y_i$$

is precisely the above formula for $\mathfrak{G}_w(\mathbf{x}, \mathbf{y})$. Similarly, (4.2) is the known formula for $\mathfrak{S}_w(\mathbf{x}, \mathbf{y})$ after the substitution

$$t_{ij} \mapsto x_j - y_i.$$

Hence (4.3) and (4.4) hold.

Note that the substitution $t_{ij} \mapsto x_j/y_i$ is precisely the map on K-polynomials induced by the inclusion of the usual action (with relabeled weights) into the rescaling action, and

the substitution $t_{ij} \mapsto x_j - y_i$ is the equivalent map for multidegrees. Thus the claim of the final sentence follows from Theorem 2.1, Theorem 3.2, and Proposition 4.1 combined. $\square$

While Definition–Theorem 4.3 exploits the usual action on $\mathcal{N}_{w_0 \star w_0, \hat{w}}$ for the double Schubert and Grothendieck polynomials, the usual action on arbitrary $\mathcal{N}_{v,w}$ can be used to geometrically explain the specialization formula [BucRim04] for double Grothendieck polynomials. (We emphasize that the following result and its proof hold even if we define Grothendieck polynomials as traditionally done [LasSch82b].)

Theorem 4.5. *We have the equalities:*

$$(4.5) \quad \mathfrak{G}_{w_0 w}(t_{v(1)}, \dots, t_{v(n)}; t_n, t_{n-1}, \dots, t_1) = \mathcal{K}(\mathcal{N}_{v,w}, t_{ij} \mapsto t_{v(j)}/t_{n-i+1}) \\ = \mathfrak{G}_{v,w}(t_{ij} \mapsto t_{v(j)}/t_{n-i+1}),$$

and

$$(4.6) \quad \mathfrak{S}_{w_0 w}(t_{v(1)}, \dots, t_{v(n)}; t_n, t_{n-1}, \dots, t_1) = \mathcal{C}(\mathcal{N}_{v,w}, t_{ij} \mapsto t_{v(j)} - t_{n-i+1}) \\ = \mathfrak{S}_{v,w}(t_{ij} \mapsto t_{v(j)} - t_{n-i+1}).$$

The equality of the first and third polynomials in each of (4.5) and (4.6) was obtained (up to differences in conventions) by [BucRim04], who furthermore ask for a geometric explanation. We respond to that question by showing that both are in fact equal to an equivariant (K-theory) class of $\mathcal{N}_{v,w}$. Our proof utilizes our Gröbner basis result, Theorem 2.1.

Proof of Theorem 4.5: The injection $\{e_v\} \hookrightarrow \text{Flags}(\mathbb{C}^n)$ induces a map

$$K_T(\text{Flags}(\mathbb{C}^n)) \rightarrow K_T(e_v) \\ [\mathcal{O}_{X_w}]_T \mapsto [\mathcal{O}_{X_w}]_T|_{e_v}.$$

Here T is the torus $(\mathbb{C}^*)^n$ and the torus action is the usual action from Section 4.1.

Since $v\Omega_{\text{id}}^\circ$ T -equivariantly contracts to e_v , and the isomorphism of Equation 2.1 is T -equivariant, we can identify the class

$$[\mathcal{N}_{v,w}] \in K_T(\Omega_v^\circ)$$

with

$$[\mathcal{O}_{X_w}]_T|_{e_v} \in K_T(e_v).$$

Choosing the usual coordinates for the weight space of T , the class in $K_T(\Omega_v^\circ)$ of any subscheme (or, in general $\mathcal{O}$ -module) over Ω_v° is given by its K -polynomial. Therefore, we can make the identifications

$$[\mathcal{N}_{v,w}] = [\mathcal{O}_{X_w}]_T|_{e_v} = \mathcal{K}(\mathcal{N}_{v,w}, t_{ij} \mapsto t_{v(j)}/t_{n-i+1}),$$

by the discussion of Section 4.1.

On the other hand, it is a folklore theorem that

$$[\mathcal{O}_{X_w}]_T|_{e_v} = \mathfrak{G}_{w_0 w}(x_j \mapsto t_{v(j)}, y_i \mapsto t_{w_0(i)}).$$

While we could not find an explicit proof for this statement in the literature, R. Goldin [Gol01] gave a proof for the equivalent statement for equivariant cohomology; her proof can be seen to extend to equivariant K -theory, with the appropriate technical modifications.

Alternatively, one can also construct a similar proof, substituting homological algebra for geometry, as sketched below. First we impose the torus action on M_n so that the variables in $\mathbf{z}$ have the weights they would have in $\mathbf{z}^{(v)}$; this means giving the variable z_{ij} the weight $t_{v(j)}/t_{w_0(i)}$ rather than the weight x_j/y_i . Now we consider the restriction map

$$K_T(\mathbb{C}[\mathbf{z}]) \rightarrow K_T(\mathbb{C}[\mathbf{z}^{(v)}]).$$

Since it is equivariantly contractible to the identity map from a point to a point, it is the identity map on K -polynomials. Furthermore, since this is a map of affine schemes, left-derived pullback is simply Tor , so for any $\mathbb{C}[\mathbf{z}]$ -module Y , the class $[Y] \in K_T(\mathbb{C}[\mathbf{z}])$ is mapped to

$$\sum_i (-1)^i [\text{Tor}_i(Y, \mathbb{C}[\mathbf{z}^{(v)}])].$$

Equivalently, if one wants to use only algebraic arguments, this can be seen from the calculation of Tor using a free resolution of Y and the calculation of K -polynomials from a free resolution (or indeed any exact sequence).

Now $\text{Tor}_0(\mathbb{C}[\mathbf{z}]/I_{w_0w}, \mathbb{C}[\mathbf{z}^{(v)}])$ is simply the coordinate ring of $\mathcal{N}_{v,w}$. Furthermore, the ideal defining $\mathbb{C}[\mathbf{z}^{(v)}]$ (defined in Section 2.2) is generated by elements of $\mathbb{C}[\mathbf{z}]$ which are a regular sequence on the coordinate ring $\mathbb{C}[\mathbf{z}]/I_{w_0w}$ of the matrix Schubert variety $\bar{X}_{w_0w}$. (It can be easily seen that they are part of a system of parameters, since the ideal is generated by $\binom{n+1}{2} + \ell(v)$ elements (as $\binom{n}{2} + \ell(v)$ entries are set to 0 and n entries to 1) and the codimension of $\mathcal{N}_{v,w}$ in $\bar{X}_{w_0w}$ is also $\binom{n+1}{2} - \ell(v)$. Since $\bar{X}_{w_0w}$ is known to be Cohen–Macaulay [Ful92], any part of a system of parameters is also a regular sequence.) Therefore, by standard facts (see for example [BruHer98, Prop. 1.6.9, Thm. 1.6.17b]) in the theory of regular sequences, $\text{Tor}_i(\mathbb{C}[\mathbf{z}]/I_w, \mathbb{C}[\mathbf{z}^{(v)}]) = 0$ for $i > 0$.

Therefore, the K_T -class for $\bar{X}_w$ (with respect to the action which restricts to the usual action on $\mathbb{C}[\mathbf{z}^{(v)}]$) restricts to the K_T class for $\mathcal{N}_{v,w}$. Since this restriction map is the identity on K -polynomials, we must have that

$$[\mathcal{N}_{v,w}] (= [\mathcal{O}_{X_w}]_T |_{e_v}) = \mathfrak{S}_{w_0w}(x_j \mapsto t_{v(j)}, y_i \mapsto t_{w_0(i)}),$$

as desired.

Thus we obtain the first equality of (4.5). Since $[\mathcal{O}_{\mathcal{N}_{v,w}}]$, the K -polynomial of $\mathcal{N}_{v,w}$, is preserved under Gröbner degeneration, the second equality of (4.5) follows from Theorem 2.1, Theorem 3.2 and Proposition 4.1 combined.

The proof of (4.6) is similar (although in this case we can directly use Goldin’s result). $\square$

Example 4.6. For simplicity, let us only illustrate the Schubert polynomial assertions (4.6) of Theorem 4.5. We continue Example 4.2 where $v = 31452$ and $w = 53142$ (and hence $w_0w = 13524$). Now, from Example 4.4, we have that

$$\begin{aligned} \mathfrak{S}_{w_0w}(t_{v(1)}, \dots, t_{v(n)}; t_n, \dots, t_1) &= \mathfrak{S}_{13524}(t_3, t_1, t_4, t_5, t_2; t_5, t_4, t_3, t_2, t_1) = \\ &= (t_1 - t_3)(t_3 - t_4)(t_1 - t_4) + (t_1 - t_3)(t_3 - t_4)(t_4 - t_5) + (t_1 - t_3)(t_1 - t_5)(t_4 - t_5) \\ &\quad + (t_3 - t_2)(t_3 - t_4)(t_4 - t_5) + (t_3 - t_2)(t_3 - t_4)(t_1 - t_4) + (t_3 - t_2)(t_1 - t_5)(t_4 - t_5) \end{aligned}$$

On the other hand, from Example 4.2, we see that

$$\begin{aligned} \mathfrak{S}_{v,w}(t_{ij} \mapsto t_{v(j)} - t_{n-i+1}) &= \mathfrak{S}_{31452,53142}(t_{ij} \mapsto t_{v(j)} - t_{n-i+1}) = \\ &= (t_1 - t_2)(t_3 - t_4)(t_1 - t_4) + (t_1 - t_2)(t_3 - t_4)(t_4 - t_5) + (t_1 - t_2)(t_1 - t_5)(t_4 - t_5). \end{aligned}$$

Now, Theorem 4.5 asserts that these two polynomials in the t_i 's are equal, which the reader can check by direct computation. However, this equality is not obvious *a priori*. $\square$

4.4. A rationale for unspecializing. The standard definition for Grothendieck and Schubert polynomials is in terms of (isobaric) divided difference operators. Our rationale for presenting this nonstandard (and highly ahistorical) definition through “unspecialization” is as follows. What [KnuMil05] taught us is that the Grothendieck and Schubert polynomials, being presented as polynomials in the x_i, y_j variables is already “biased” as a equivariant (K-theory) class of a matrix Schubert variety for the special torus action. However, from this point of view, the specialization of these polynomials examined in Theorem 4.5 appears geometrically unnatural and even combinatorially mysterious. Our approach seeks to emphasize that rather than viewing the latter as a specialization of the former, one should think that the two are specializations of two different though related unspecialized Grothendieck polynomial. Moreover, these specializations are geometrically natural, since they arise as explained in Section 4.2 from a restriction from the larger rescaling torus action to the smaller usual torus action.

5. MULTIPLICITIES OF SCHUBERT VARIETIES

We take this opportunity to relate our work to the problem of finding positive formulas for multiplicities of Schubert varieties.

The **multiplicity** of a point p in a scheme X , denoted $\text{mult}_p(X)$ is defined as the degree of the projective tangent cone $\text{Proj}(\text{gr}_{\mathfrak{m}_p} \mathcal{O}_{X_p})$ as a subvariety of the projective tangent space $\text{Proj}(\text{Sym}^* \mathfrak{m}_p / \mathfrak{m}_p^2)$, where $(\mathcal{O}_{X_p}, \mathfrak{m}_p)$ is the local ring associated to $p \in X$. Equivalently, if the Hilbert–Samuel polynomial of $\mathcal{O}_{X_p}$ is $a_d x^d + a_{d-1} x^{d-1} + \dots + a_0$ then

$$\text{mult}_p(X) = d! a_d.$$

The following open problem has been of interest:

Problem 5.1. *Give an explicit, nonrecursive, positive combinatorial rule to compute $\text{mult}_{e_v}(X_w)$, for each $(v, w) \in S_n \times S_n$.*

This problem remains open for (most cases of) generalized flag varieties G/P , and even for the case of $\text{Flags}(\mathbb{C}^n)$, which is our present focus. However, for minuscule G/P , a recursive formula was given by Lakshmibai and Weyman [LakWey90]. For the special case of the Grassmannian $\text{Gr}(k, \mathbb{C}^n)$ of k -dimensional planes in $\mathbb{C}^n$, several closed formulas have been given [RosZel01, Kra01, KreLak04], and similar formulas were recently given for the symplectic Grassmannian [GhoRag06] and the orthogonal Grassmannian [RagUpa07] (both in the case of maximal isotropic subspaces).

5.1. Homogeneity and parabolic moving. Let us now describe two facts and a conjecture, which allow us to positively compute multiplicities in many cases. The first gives a combinatorial rule whenever the Kazhdan–Lusztig ideal is homogeneous. The proof was suggested to us by A. Knutson in a private communication of his combinatorial rule for multiplicity problems (v, w) which holds in any type whenever w_0v is λ -cominusculer (for some weight λ). In particular w_0v is λ -cominusculer in type A if it is 321-avoiding. See [Ste01, Proposition 2.1] and Section 4.1 for details.

Fact 5.1 (Homogeneity). *Suppose $v, w \in S_n$ and $I_{v,w}$ is homogeneous with respect to the standard grading that assigns $\deg(z_{ij}) = 1$. Then $\text{mult}_{e_v}(X_w)$ equals the number of facets of $\Delta_{v,w}$, or equivalently the number of reduced pipe dreams for w on $D(v)$.*

Proof. The value of $\text{mult}_{e_v}(X_w)$ equals the degree of the initial ideal of $I_{v,w}$ with respect to any term order $\prec'$ that always picks out a lowest degree term as its leading term. However, if $I_{v,w}$ is already homogeneous with respect to the standard grading, then one can use $\prec' := \prec$, and thus the result follows from Theorems 2.1 and 3.2. $\square$

Ideally, one would have a simple combinatorial characterization for when $I_{v,w}$ is homogeneous with respect to the standard grading (see [WooYon08, Problem 5.5]). At present, we do not know how to solve even the presumably simpler problem of determining when the defining (or essential) minors are homogeneous.

For the purposes of computing multiplicity in general, we would need, as stated in the proof above, a Gröbner basis under any term order that picks out a lowest degree term. The defining determinants are not a Gröbner basis in general for any such term orders we have tried. However, as we shall see below, it suffices to solve only a subset of these problems.

To see this, let us now recall another well-known trick. Let

$$\mathcal{T} = \{s_i = (i \leftrightarrow i+1) \mid s_i w < w\}.$$

These are known as the **left descents** of w . Similarly let

$$\mathcal{T}' = \{s_i = (i \leftrightarrow i+1) \mid w s_i < w\}$$

be the set of **right descents** of w . In general, given a Schubert variety X_w , the parabolic subgroup

$$P_{\mathcal{T}} \subset G$$

generated by B and the transpositions in $\mathcal{T}$ acts on it by left multiplication. In particular, if $s_i w < w$, then this action induces an isomorphism of a local neighbourhood of e_v in X_w with a local neighbourhood of $e_{s_i v}$ in X_w , thus preserving all local properties at these points. Moreover, since local properties are preserved under inverse (as $\mathcal{N}_{v,w}$ is isomorphic to $\mathcal{N}_{v^{-1}, w^{-1}}$), we also have that, if $w s_i < w$, then a similar statement holds for e_v and $e_{v s_i}$ in X_w . Summarizing, one can compute invariants of e_v in X_w from $e_{v'}$ in X_w whenever v and v' are in the same double coset $S_{\mathcal{T}} v S_{\mathcal{T}'}$ in S_n . Here $S_{\mathcal{T}}$ and $S_{\mathcal{T}'}$ denote respectively the subgroups of S_n generated by the simple transpositions in $\mathcal{T}$ and $\mathcal{T}'$.

Applying the above trick to computing multiplicity, we have:

Fact 5.2 (Parabolic moving). *If $s_i w < w$, then $\text{mult}_{e_v}(X_w) = \text{mult}_{e_{s_i v}}(X_w)$. Similarly, if $w s_i < w$, then $\text{mult}_{e_{v s_i}}(X_w) = \text{mult}_{e_v}(X_w)$.*

Combining Facts 5.1 and 5.2, one can hope to positively compute $\text{mult}_{e_v}(X_w)$ by using parabolic moving to instead calculate $\text{mult}_{e_{v'}}(X_w)$ where

$$v' \in S_{\mathcal{T}} v S_{\mathcal{T}'}$$

and $I_{v',w}$ is standardly homogeneous.

Actually, we expect conjecturally that there is a particular “good” $v' \in S_{\mathcal{T}} v S_{\mathcal{T}'}$ to use. Define

$$v' = v_{\max}$$

to be **parabolically maximal** if it is maximal (in Bruhat order) in its double coset $S_{\mathcal{T}} v S_{\mathcal{T}'}$. Each double coset has a unique maximal element. Combinatorially, if v' is parabolically maximal (for some v) if its left and right descent sets contain those of w . Moreover given v , we can find $v_{\max}$ by first rearranging in decreasing order the entries of v with numbers corresponding to segments of consecutive generators of $\mathcal{T}$, then rearranging the resulting permutation so that entries in positions corresponding to segments of consecutive generators of $\mathcal{T}'$ are in decreasing order.

Example 5.2. Let $v = 316298475 \leq w = 896354721$. Then the left descents and right descents of w are given respectively by

$$\mathcal{T} = \{s_1, s_2, s_4, s_5, s_7\} \text{ and } \mathcal{T}' = \{s_2, s_3, s_5, s_7, s_8\}.$$

In order to obtain $v_{\max}$, the elements s_1 and s_2 of $\mathcal{T}$ indicate that one should rearrange the labels 1, 2, 3 in v in decreasing order, whereas the elements s_4 and s_5 of $\mathcal{T}$ indicate that one should then rearrange the labels 4, 5, 6 in v in decreasing order, and s_7 indicates that 8 should be put before 7. Doing this, one obtains $v \mapsto 326198574$. Now, similarly, the elements s_2 and s_3 of $\mathcal{T}'$ tell us to rearrange the positions 2, 3, 4 of 326198574, and so on. This process then terminates with $v_{\max} = 362198754$. $\square$

Our discussion above shows that to solve all multiplicity problems, it suffices to solve the parabolically maximal ones. The following asserts that it suffices to check the homogeneity of $I_{v_{\max},w}$ if one wishes to know if Facts 5.1 and 5.2 suffice to compute the multiplicity of e_v on X_w .

Conjecture 5.3 (Parabolic maximality). *Suppose the Kazhdan–Lusztig ideal $I_{v,w}$ is standardly homogeneous. If $ws_i < w$ but $vs_i > v$, then $I_{vs_i,w}$ is standardly homogeneous, and similarly, if $s_i w < w$ but $s_i v > v$, then $I_{s_i v,w}$ is standardly homogeneous. Therefore, $I_{v_{\max},w}$ is standardly homogeneous.*

The final sentence of Conjecture 5.3 clearly follows by induction using the second sentence. It seems plausible that one can deduce the second sentence using similar analysis as in our proof of Proposition 6.15 in Section 6.2, although we do not pursue this here.

5.2. Computational results and Monte Carlo simulation. Together Facts 5.1, 5.2 and Conjecture 5.3 provide a useful means to solve multiplicity problems. We can use the symbolic algebra software Macaulay 2 to computationally check whether $I_{v,w}$ is standardly homogeneous, by first applying `trim` to the set of defining (or essential) minors of $I_{v,w}$ and then using the function `isHomogeneous`. Testing on the set

$$\Gamma_n := \{(v, w) \in S_n \times S_n \mid v < w \text{ in Bruhat order}\},$$

we found by exhaustive search that, for $n = 5$, 74% of all problems fall to Fact 5.1 alone. However, not suprisingly, the success percentage falls off quickly.

On the other hand, if we also use Fact 5.2 and consistently replace $(v, w) \in \Gamma_n$ with $(v_{\max}, w) \in \Gamma_n$, the success percentage increases rather substantially. By exhaustive search, all problems for $n \leq 4$ are solved this way while 98.5% of the 3871 problems are solved for $n = 5$. Moreover, Monte Carlo simulation estimates are summarized in the following table:

n	6	7	8	9	10
success %	94	86	73	62	46

TABLE 1. Estimates of success percentage with 2,000 Monte Carlo trials, using Facts 5.1 and 5.2.

We found it encouraging that such simple tricks allow one to cover such a large fraction of all multiplicity problems for even up to $n = 10$.

Let us collect a few more computationally determined facts:

Fact 5.3. (I) $I_{v_{\max}, w}$ need not be standardly homogeneous, one example is

$$I_{13425,34512} = \langle z_{11}, z_{12}, z_{21}, z_{13}z_{22}z_{31} - z_{14}z_{41} \rangle$$

(Therefore, Conjecture 5.3, even if true, would not solve all multiplicity problems).

(II) $I_{v, w}$ may be standardly homogeneous even if $w_0 v$ is not 321-avoiding (and therefore not λ -cominusculc for any λ); for example

$$I_{45213,54231} = \langle z_{33} \rangle.$$

(III) $I_{v, w}$ might not be standardly homogeneous even if $I_{v_{\max}, w}$ is. (Hence the converse to Conjecture 5.3 is false.) For example,

$$I_{v, w} = I_{31524,43512} = \langle z_{11}, z_{12}, z_{24}z_{42} - z_{22} \rangle$$

while

$$I_{v_{\max}, w} = I_{41532,43512} = \langle z_{11}, z_{12}, z_{42} \rangle.$$

It is worthwhile to mention that, as with all checks on Γ_n , the computational demands are large for $n \geq 6$. On the other hand, it is not difficult to Bernoulli sample a pair $(v, w) \in \Gamma_n$ uniformly at random. One can independently and uniformly pick two permutations $\sigma, \rho \in S_n$, and reject until either $\sigma \leq \rho$ or $\rho \leq \sigma$. In the first case, one returns $(u, v) = (\sigma, \rho)$ while in the latter case one returns $(u, v) = (\rho, \sigma)$.

In our experience, this approach allows one to practically estimate success probabilities for n beyond the reach of exhaustive search, in the sense that the true bottleneck in computation comes from the Gröbner basis computations. In particular, Conjecture 5.3 is endorsed up to $n \leq 10$ using this method.

We also remark that rigorous analysis of the likelihood of picking a Bruhat comparable pair from $S_n \times S_n$ was performed recently by A. Hammett and B. Pittel [HamPit08]. They bound this probability by

$$c_1(0.708^n) \leq \mathbb{P}[u \leq v \text{ or } v \leq u | (u, v) \in S_n \times S_n] \leq c_2 \frac{1}{n^2},$$

for universal constants $c_1, c_2 > 0$. Moreover they conjecture this probability is about $n^{-3/2}$. If their conjecture is true, then one would expect the Bernoulli sampling algorithm to terminate quickly, in about $O(n^{3/2})$ trials, which agrees with our experience.

The above computations support the idea that Monte Carlo simulation is a useful resource when studying algebraic combinatorics and computational commutative algebra such as that present in [WooYon08]. In that work, one needs to sample elements of Γ_n satisfying “interval pattern avoidance conditions”. This motivates the need for more sophisticated (and efficient) sampling algorithms (via methods such as Markov Chain Monte Carlo or importance sampling); further discussion may appear elsewhere.

5.3. Formulas in the Grassmannian case. Now suppose $w \in S_n$ is co-Grassmannian, meaning that it has a unique ascent $w(k) < w(k+1)$ (or, equivalently, that there is at most one simple transposition s_k with the property that $ws_k > w$). Let us consider the multiplicity problem in only this case. In general, if $v \leq w$, it is not true that v is also **co-Grassmannian**. However, observe that we can always replace v by $v_{\max}$, which is then (by the discussion of Section 5.1), co-Grassmannian, with its unique ascent also in position k . This reduces the problem to computing multiplicities on Grassmannians, a problem previously considered in [LakWey90, RosZel01, Kra01]. In summary these results provide determinantal and tableau based formulas for the multiplicity. Our goal here is to provide a (mildly) more general, simpler, formula (being valid for all $v \leq w$ and not only co-Grassmannian $v \leq w$), together with a new conceptual explanation for the appearance of these formulas, using the results of this paper.

Let us therefore assume unless otherwise stated that $v \leq w$ are both co-Grassmannian with the same ascent position k . The co-Grassmannian assumption on v allows us to easily check that the defining generators of $I_{v,w}$ are homogeneous with respect to the standard grading. In fact, homogeneity also follows from w_0v being λ -cominuscul.

Consequently, Fact 5.1 guarantees that the multiplicity of e_v in X_w is the number of reduced pipe dreams on $D(v)$ for w_0w . Note that w_0w is a **Grassmannian permutation**, meaning one with a unique descent, in this case at position k . Moreover, under the present assumptions one can establish a bijection between reduced pipe dreams for w_0w in $D(v)$ and “flagged” semistandard tableaux, as we explain below.

First we need some standard facts about the combinatorics of co-Grassmannian permutations. To each such permutation w with its unique ascent at position k , we associate a partition

$$\lambda(w) = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k \geq 0)$$

by setting

$$\lambda_{k-i+1} = n - w(i) + 1 - i.$$

Hence, for example, if $w = 975386421$ then $\lambda(w) = (4 \geq 2 \geq 1 \geq 0)$. If v and w are two co-Grassmannian permutations both with ascents at k , then $\lambda(v) \supseteq \lambda(w)$ if and only if $v \leq w$. Note under these conventions $|\lambda| = \sum_i \lambda_i$ is the codimension of X_w in $\text{Flags}(\mathbb{C}^n)$, which $\binom{n}{2} - \ell(w)$ or equivalently the number of non-inversions in w .

The boxes of $\text{flatten}(D(v))$ form the shape $\lambda(v)$ rotated 180 degrees and conjugated (transposed). Now consider the flattened pipe dreams inside $\text{flatten}(D(v))$, as in Section 3.3. The co-Grassmannian assumption on w (and hence Grassmannian assumption on w_0w) implies that we can produce each reduced pipe dream for w_0w by the following procedure, which we describe in terms of pipe dreams drawn on $\text{flatten}(D(v))$ rather

than on $D(v)$. One can easily recover reduced pipe dreams in $\text{RedPipes}(v, w_0 w)$ by unflattening.

Start with the **starting pipe dream**, which is the unique pipe dream whose $+$'s form the shape $\lambda(w)$ rotated 180 degrees, conjugated, and placed in the lower right hand corner of $\text{flatten}(D(v))$. Then, locally, one can make the transformation

$$(5.1) \quad \begin{array}{cc} \cdot & \cdot \\ \cdot & + \end{array} \mapsto \begin{array}{cc} + & \cdot \\ \cdot & \cdot \end{array}$$

where each 2×2 configuration describes a subsquare of $\text{flatten}(D(v))$, and the “ $\cdot$ ” refers to a square of $\text{flatten}(D(v))$ without a $+$. Such a transformation will produce another reduced pipe dream for $w_0 w$, and one can generate any reduced pipe dream for $w_0 w$ which fits inside $\text{flatten}(D(v))$ by some sequence of such transformations from the starting pipe dream.

We now associate a semistandard Young tableau of shape $\lambda(w)$ to each reduced pipe dream for $w_0 w$ on $\text{flatten}(D(v))$. We associate to the starting pipe dream the super-semistandard **starting tableau** of shape $\lambda(w)'$, defined to have label m in every box of row m . More generally, each $+$ in the starting tableau is in obvious bijection with a box of $\lambda(w)$, and following the local transformation (5.1) allows one to coherently associate each $+$ of any pipe dream to a box of $\lambda(w)$, namely the box associated to the $+$ in the starting tableau that it came from. Now if a $+$ is in the i -th column of $\text{flatten}(D(v))$, counting from the *right* (and starting with the first column having a box), then we put an i in the corresponding box of $\lambda(w)$. The resulting tableau can be seen (by induction) to be semi-standard. These conclusions essentially follow from the analysis of [KnuMilYon05] together with Section 3.3.

For example, if $\lambda(v) = (5, 4, 4, 2)$ and $\lambda(w) = (4, 2, 1, 0)$ we have that the starting pipe dream (after rotating 180-degrees) and the starting tableau are (after flattening, rotating and conjugating):

$$(5.2) \quad \begin{array}{|c|c|c|c|c|} \hline + & + & + & + & \\ \hline + & + & & & \\ \hline + & & & & \\ \hline & & & & \\ \hline \end{array} \leftrightarrow \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 1 \\ \hline 2 & 2 & & \\ \hline 3 & & & \\ \hline \end{array},$$

and all others are obtained by the local moves (5.1), rotated, which look like

$$(5.3) \quad \begin{array}{cc} + & \cdot \\ \cdot & \cdot \end{array} \mapsto \begin{array}{cc} \cdot & \cdot \\ \cdot & + \end{array}$$

Not every semistandard tableau of shape $\lambda(w)$ can be obtained this way. The maximum entry of row m of such a tableau T is bounded from above by how far south the rightmost $+$ in the m -th row of the starting pipe dream can travel and remain inside $\lambda(v)$. Thus in the above example, the possible semistandard tableau are of shape $\lambda(w) = (4, 2, 1)$ such that the entries in the first, second and third rows respectively are bounded by 1, 3, and 4. In general, this is given by b_m , which is the row at the bottom of the largest square that contains the right most box of $\lambda(w)_m$ as its northwest corner and is contained inside $\lambda(v)$. In symbols:


$$b_m = \max_i \{ \lambda(v)_i \geq \lambda(w)_m + i - m \}.$$

Clearly the sequence $b = (b_1, b_2, \dots, b_k)$ is weakly increasing.


Example 5.4. Let $v = 743198652 \leq w = 975286431$. Then

$$\lambda(v) = (5, 4, 4, 2) \supseteq \lambda(w) = (4, 2, 1, 0).$$

The diagram $D(v)$ and its canonical labeling are depicted below:




After flattening $D(v)$ we obtain the shape $(4, 4, 3, 3, 1)$ as read from bottom to top. This is the conjugate shape of $\lambda(v)$. The starting pipe dream is given by



The reader can check that the associated reduced word for this starting pipe dream is $s_7 s_6 s_4 s_5 s_2 s_3 s_4 = w_0 w = 135824679$.

After conjugating and rotating, the pipe dreams are precisely those which can be obtained via a sequence of local moves from (5.3) the starting pipe dream depicted in (5.2). These are then in natural bijection with the semistandard tableaux with row bounds $(1, 3, 4)$:



Hence $\text{mult}_v(X_w) = 5$. □

The weight generating series

$$\sum_T \mathbf{x}^{\text{wt}(T)}$$

where the sum runs over all semistandard tableaux of shape λ with row entries **flagged** (bounded) by a vector $\mathbf{d}$ is called the **flagged Schur polynomial**. A standard Gessel–Viennot type argument shows that

$$\sum_{\mathbf{T}} \mathbf{x}^{\text{wt}(\mathbf{T})} = \det(h_{\lambda_i - i + j}(x_1, \dots, x_{b_i})),$$

where $h_d(x_1, \dots, x_b)$ is the complete homogeneous symmetric function of degree d in the variables $x_1, \dots, x_b$. See [Man01, Cor 2.6.3] for details.

By setting $x_1 = x_2 = \dots = 1$ into this formula, we obtain a formula for the multiplicity as a determinant of a matrix with binomial coefficient entries.

The above discussion therefore proves the following theorem:

Theorem 5.5. *Let $w \in S_n$ be a co-Grassmannian permutation with unique ascent at position k . Then if $v \leq w$, we have that $v_{\max}$ is co-Grassmannian with unique ascent at position k , and $\text{mult}_{e_v}(X_w)$ equals the number of semistandard flagged Young tableau of shape $\lambda = \lambda(w)$ flagged by the vector $\mathbf{b}$ given by*

$$b_m = \max_i \{ \lambda(v_{\max})_i \geq \lambda(w)_m + i - m \}.$$

In addition, we have the determinantal formula

$$\text{mult}_{e_v}(X_w) = \det \left(\binom{b_i + \lambda_i - i + j - 1}{\lambda_i - i + j} \right)_{1 \leq i, j \leq \ell(\lambda)},$$

where $\ell(\lambda)$ is the number of nonzero parts of λ .

Example 5.6. Continuing the previous example, we have

$$\text{mult}_{e_v}(X_w) = \begin{pmatrix} \binom{b_1 + \lambda_1 - 1}{\lambda_1} & \binom{b_1 + \lambda_1}{\lambda_1 + 1} & \binom{b_1 + \lambda_1 + 1}{\lambda_1 + 2} \\ \binom{b_2 + \lambda_2 - 2}{\lambda_2 - 1} & \binom{b_2 + \lambda_2 - 1}{\lambda_2} & \binom{b_2 + \lambda_2}{\lambda_2 + 1} \\ \binom{b_3 + \lambda_3 - 3}{\lambda_3 - 2} & \binom{b_3 + \lambda_3 - 2}{\lambda_3 - 1} & \binom{b_3 + \lambda_3 - 1}{\lambda_3} \end{pmatrix} = \begin{pmatrix} \binom{4}{4} & \binom{5}{4} & \binom{6}{5} \\ \binom{3}{3} & \binom{4}{3} & \binom{5}{4} \\ \binom{2}{2} & \binom{3}{3} & \binom{4}{4} \end{pmatrix} = 5,$$

in agreement with our previous computation. $\square$

Although our formula is also a determinant of a matrix of binomial coefficients, it is different from the ones given in [LakWey90] and [RosZel01]. Presumably it would not be difficult to show the formulas are equivalent through a succession of determinantal identities. Our tableau rule is also different than the one given by [Kra01] to explain the positivity and equivalence of these two determinantal expressions; his rule instead counts semistandard tableaux of an irregular shape satisfying certain column and row bounds. The intermediate pipe dream arguments we use, as we have said, are closely related to [KnuMilYon05]. However, they also appear in later work, specifically of V. Kreiman [Kre08] and T. Ikeda and H. Naruse [IkeNar07], for reasons similar to ours. The rather trivial distinction is that these authors focus on T -fixed points on the Grassmannian itself rather than on Schubert varieties in the flag variety indexed by co-Grassmannian permutations as we do here.

Our proof gives a Gröbner geometry explanation of the appearance of tableaux: the multiplicity is the degree of the Kazhdan–Lusztig ideal $I_{v,w}$, which can via Theorems 2.1 and 3.2 be thought of geometrically as the number of components of $\text{in}_{\prec} I_{v,w}$ and combinatorially as the number of pipe dreams. These pipe dreams are in this case transparently in bijection with flagged semistandard tableaux.

Perhaps notable is the appearance of *flagged* tableaux in our formulas for the multiplicity. Flagged tableaux and flagged Schur functions most often appear in the combinatorics of co-vexillary permutations and their Schubert polynomials. While co-Grassmannian permutations are co-vexillary, our proof does not extend in general to cases where w or v is co-vexillary. Algebraically, this amounts to the fact that the Kazhdan–Lusztig ideal is no longer homogeneous with respect to the standard grading (even if one replaces v by $v_{\max}$). Instead, a more refined degeneration argument is needed; the second author is presently investigating this problem with L. Li.

The argument we use should work more generally to give formulas for $X_w \subseteq G/B$ when $w_0 w$ is λ -cominuscule. In particular, one should be able to obtain determinantal formulas for multiplicities of Schubert varieties of G/P where P is a co-minuscule maximal parabolic.

6. PROOF OF THEOREMS 2.1 AND 3.2

Let $R = \mathbb{C}[z^{(v)}]$ and suppose

$$I_{v,w} := \langle m_1, \dots, m_N \rangle \subseteq R$$

where $m_1, \dots, m_N$ (for some N) are the essential minors, as explained in Section 1.2 (using (3.1 and (3.2)). Let

$$(6.1) \quad J_{v,w} := \langle LT_{\prec}(m_1), \dots, LT_{\prec}(m_N) \rangle \subseteq \text{in}_{\prec} I_{v,w}$$

be the ideal generated by the leading terms of the essential minors, with respect to the term order $\prec$ defined in Section 1.3. By definition, the containment is an equality if and only if $\{m_1, \dots, m_N\}$ is a **Gröbner basis** with respect to $\prec$.

The key technical step for our proof of Theorems 2.1 and 3.2 is the following.

Theorem 6.1. *If a pipe dream $\mathcal{P}$ does not label a face of $\Delta_{v,w}$, then the corresponding monomial $z^{\mathcal{P}}$ in R is divisible by one of the leading terms $LT_{\prec}(m_1), \dots, LT_{\prec}(m_N)$ of an essential minor of $J_{v,w}$.*

The converse also holds, and follows from Theorem 2.1, but we will not need this.

We also need the following theorem.

Theorem 6.2. *Let $R/K_{v,w}$ be the Reisner–Stanley ring of $\Delta_{v,w}$. Then*

$$\mathcal{K}(R/K_{v,w}, \mathbf{t}) = \mathcal{K}(R/I_{v,w}, \mathbf{t}),$$

where the K -polynomials are calculated relative to the grading given by the usual action defined in Section 4.1.

Delaying the proof of Theorems 6.1 and 6.2, we are now ready to give the:

Conclusion of the proof of Theorem 2.1: Let

$$K_{v,w} = \text{the Reisner–Stanley ideal of the pipe complex } \Delta_{v,w}.$$

Theorem 6.1 implies that if $\mathcal{P}$ is a nonface of $\Delta_{v,w}$ then $\mathcal{P}$ is in $J_{v,w}$. Hence

$$(6.2) \quad K_{v,w} \subseteq J_{v,w} \subseteq \text{in}_{\prec} I_{v,w},$$

where the latter containment reiterates (6.1).

So by (6.2) we have surjections

$$R/K_{v,w} \twoheadrightarrow R/J_{v,w} \twoheadrightarrow R/\text{in}_{\prec} I_{v,w}.$$

Theorem 6.2 states that

$$\mathcal{K}(R/K_{v,w}, \mathbf{t}) = \mathcal{K}(R/I_{v,w}, \mathbf{t}).$$

Hence the above containments are actually equalities, and

$$K_{v,w} = J_{v,w} = \text{in}_{\prec} I_{v,w}.$$

Thus the essential minors of $I_{v,w}$ are Gröbner with respect to $\prec$. Moreover $\Delta_{v,w}$ is the Reisner–Stanley complex of the initial ideal.

The above argument only proves the theorem when $\mathbb{k} = \mathbb{C}$, since we have used facts about Schubert varieties that are proved only for that case. (See, specifically, the proof of Theorem 6.2.) However, the general case follows since all coefficients of terms in the essential minors are ± 1 . $\square$

Conclusion of the proof of Theorem 3.2: In the proof of Theorem 2.1 we saw that $\Delta_{v,w}$ is the Reisner–Stanley complex of the initial ideal $\text{in}_{\prec} I_{v,w}$. By Proposition 3.3, $\Delta_{v,w}$ is homeomorphic to a ball or sphere. In particular it is equidimensional with the stated facets. Hence the prime decomposition claim follows. $\square$

In the remainder of this section, we prove Theorems 6.1 and 6.2.

6.1. Vertex decompositions of simplicial complexes. Given a simplicial complex Δ and a vertex $V \in \Delta$, the **deletion** of V is the set of the faces of Δ that do not contain V :

$$\text{del}_V(\Delta) = \{F \in \Delta : V \notin F\}.$$

The **link** of V consists of the faces in the deletion which remain faces of Δ after adding V :

$$\text{link}_V(\Delta) = \{F \in \Delta : V \notin F, F \cup \{V\} \in \Delta\},$$

and the **star** of V is:

$$\text{star}_V(\Delta) = \text{cone}_V(\text{link}_V(\Delta)).$$

Moreover, one has the decomposition

$$\Delta = \text{del}_V(\Delta) \cup \text{star}_V(\Delta).$$

A good case is when $\text{del}_V(\Delta)$ is homeomorphic to a dimension d -ball B^d and $\text{link}_V(\Delta) \cong B^{d-1}$ is on the spherical surface of $\text{star}_V(\Delta)$, whence we can deduce $\Delta \cong B^d$. This gives an inductive way of proving ballness (or similarly sphereness) of Δ by a good ordering of the vertices of Δ , and additionally implies a shelling of Δ .

The above type of reasoning was introduced in [BilPro79] and exploited in [KnuMil05] to deduce that all subword complexes, and therefore the complexes $\Delta_{v,w}$, are balls or spheres. We will use this inductive framework to prove Theorem 6.1. In [KnuMilYon05], A. Knutson, E. Miller and the second author developed a theory of “geometric vertex decompositions” by which one can inductively deduce Gröbnerness of a generating set of an ideal I by the Gröbnerness of a related generating set for a partial Gröbner degeneration I' of I . However, we will not use this theory, but rather base our induction on the following observation, whose proof is immediate from the definitions:

Lemma 6.3. *Let S be a set of vertices on any simplicial complex Δ and V a vertex of Δ .*

The set S is a nonface of Δ if and only if either:

- (1) The vertex $V \in S$, and $S \setminus \{V\}$ is a nonface of $\text{link}_V(\Delta)$.
- (2) The vertex $V \notin S$, and S is a nonface of the $\text{del}_V(\Delta)$.

Let z_{last} be the largest letter under the term order $\prec$ (which is the rightmost, then southmost variable appearing in $Z^{(v)}$). For the remainder of this paper, let V be the associated vertex of $\Delta_{v,w}$, which is the vertex labeled by the pipe dream having a cross $+$ in every position of $D(v)$ except at the position of z_{last} .

Theorem 6.4 (See [KnuMil04], proof of Theorem 2.5). *Let i be the last ascent of v and s_i the corresponding adjacent transposition.*

(A) *If i is an descent of w , then V is a cone point of $\Delta_{v,w}$, and*

$$\text{link}_V(\Delta_{v,w}) = \text{del}_V(\Delta_{v,w}) \cong \Delta_{vs_i,w}.$$

(B) *If i is an ascent of w , then*

$$\text{link}_V(\Delta_{v,w}) \cong \Delta_{vs_i,w} \quad \text{and} \quad \text{del}_V(\Delta_{v,w}) \cong \Delta_{vs_i,ws_i}.$$

In [KnuMil04], the authors find a vertex decomposition of any subword complex at the vertex associated to the last letter of Q . In view of Proposition 3.3, we can deduce Theorem 6.4 from their results (or proof). That said, to exploit the vertex decomposition in our proof of Theorem 6.1, we need to have handy the specific homeomorphisms we use in terms of the pipe complex. (See claims inside the proof below.) Explaining them amounts to a proof of Theorem 6.4 anyway.

Our proofs of Theorems 6.1 and 6.4 use the following straightforward fact:

Lemma 6.5. *Let $v \in S_n$ and i be the last ascent of v . Then the placement of boxes of $D(v)$ and $D(vs_i)$ agree in all columns except i and $i + 1$. Moreover, to obtain $D(vs_i)$ from $D(v)$, move all the boxes of $D(v)$ in column i strictly above row $v(i + 1)$ one box to the right, and also delete the box (that must appear) in position $(v(i + 1), i)$ of $D(v)$.*

Also, we need:

Lemma 6.6. *Let i be the last ascent of v . The rightmost and southmost box of $D(v)$ (which is the position of z_{last}) is in column i , and, moreover, the canonical labeling fills that box with i .*

Proof. Note that the label of a box under the canonical labeling is the “Manhattan distance” of that box from the southwest corner minus the number of dots southwest of the box. From this description, the canonical labeling is the same whether it is defined by rastering along rows or columns. The Lemma follows from the latter description. $\square$

Example 6.7. Let $v = 142653$. Then the last ascent occurs at $i = 3$. In Figure 3 we draw the diagrams for $D(v)$ and $D(vs_i)$, as an illustration of Lemma 6.5 and 6.6. $\square$

Proof of Theorem 6.4: We first prove (A). Since the facets of $\Delta_{v,w}$ are pipe dreams $\mathcal{P}$ such that $\prod \mathcal{P}$ is a reduced word for w_0w , the assertion that V is a cone point of $\Delta_{v,w}$ amounts to the following:

Claim 6.8. *No reduced pipe dream for w_0w puts a $+$ at z_{last} .*




FIGURE 3. $D(v)$ versus $D(vs_i)$

Proof of Claim 6.8: Combine Lemma 6.6 with the hypothesis that i is an ascent of w_0w . $\square$

Whenever one vertex decomposes at a cone point, the link is automatically equal to (rather than merely being a subset of) the deletion. Thus we complete (A) by proving:

Claim 6.9. *The homeomorphism between*

$$\text{del}_V(\Delta_{v,w}) = \text{link}_V(\Delta_{v,w}) \text{ and } \Delta_{vs_i,w}$$

is obtained as follows: given a pipe dream $\text{Pipe}(\mathcal{P})$ of a face $\mathcal{P} \in \text{del}_V(\Delta_{v,w})$, construct a pipe dream $\text{Pipe}(\tilde{\mathcal{P}})$ of a face $\tilde{\mathcal{P}}$ of $\Delta_{vs_i,w}$ by first deleting the $+$ in the position of z_{last} and moving each remaining $+$ in column i of $\text{Pipe}(\mathcal{P})$ one step to the right into column $i + 1$.

Proof of Claim 6.9: By Lemma 6.5, it follows that $\tilde{\mathcal{P}}$ is a pipe dream for $D(vs_i)$. Since

$$\mathcal{P} \in \text{del}_V(\Delta_{v,w}),$$

$\text{Pipe}(\mathcal{P})$ has a $+$ at the position of z_{last} . Now by Lemma 6.6, $\prod \mathcal{P}$ has an s_i at the right end of the Demazure product, and this product is by assumption equal to w_0w . But w_0w has an ascent at position i , so it follows that the same Demazure product with s_i removed still gives w_0w . This latter product is the same as $\prod \tilde{\mathcal{P}}$, so

$$\tilde{\mathcal{P}} \in \Delta_{vs_i,w}.$$

It is also clear that the map $\mathcal{P} \mapsto \tilde{\mathcal{P}}$ is injective and reversible and preserves face containment. Thus the conclusion follows. $\square$

Now we prove (B). Let us first analyze $\text{link}_V(\Delta_{v,w})$, which by definition consists of all faces $\mathcal{P} \in \Delta_{v,w}$ that

- (a) do not contain V (being in $\text{del}_V(\Delta_{v,w})$)
- (b) but satisfy $\mathcal{P} \cup V \in \Delta_{v,w}$.

Translating, (a) says that $\text{Pipe}(\mathcal{P})$ uses a $+$ in position z_{last} , whereas (b) says that removing that $+$ still gives a face of $\Delta_{v,w}$. In view of this, we have:

Claim 6.10. *The homeomorphism between*

$$\text{link}_V(\Delta_{v,w}) \text{ and } \Delta_{vs_i,w}$$

is obtained with as similar map as in Claim 6.9: given a pipe dream $\text{Pipe}(\mathcal{P})$ of a face $\mathcal{P} \in \text{link}_V(\Delta_{v,w})$, we construct a pipe dream $\text{Pipe}(\tilde{\mathcal{P}})$ of a face $\tilde{\mathcal{P}}$ of $\Delta_{vs_i,w}$ by first deleting the $+$ in the position of z_{last} and moving each remaining $+$ in column i of $\text{Pipe}(\mathcal{P})$ one step to the right, into column $i + 1$. The same map describes a homeomorphism between $\text{del}_V(\Delta_{v,w})$ and Δ_{vs_i,ws_i} .

Proof. This proof is similar to that for Claim 6.9. The key point of the link claim is that removing the $+$ in position z_{last} does not change the Demazure product. In the deletion claim, this removal of a $+$ does change the Demazure product of w , but since w has an ascent at i , the resulting Demazure product is ws_i instead. $\square$

The proof of Theorem 6.4 follows. $\square$

Example 6.11. Continuing Example 3.5, we have that $i = 3$ is an ascent of w . The vertex V is the top leftmost vertex of Figure 2 and the link is a 1-dimensional ball isomorphic to $\Delta_{vs_i, w}$, which is Example 3.4. This agrees with Theorem 6.4 and Claim 6.10. $\square$

6.2. The Kostant–Kumar recursion and the proof of Theorem 6.2. B. Kostant and S. Kumar [KosKum90] (see also [Knu08, Theorem 1]) show that the equivariant K-theory classes

$$[\mathcal{O}_{\mathcal{N}_{v,w}}] \in K_T(\Omega_v^\circ),$$

and hence the K-polynomials

$$\mathcal{K}(\mathbb{R}/I_{v,w}, t)$$

uniquely satisfy the following recursion.

Theorem 6.12. *Let $v, w \in S_n$.*

- *If $v \not\leq w$, then*

$$\mathcal{K}(\mathbb{R}/I_{v,w}, t) = 0.$$

- *If $v = w_0$, then $w = w_0$ (or we are in the previous case). Then*

$$\mathcal{K}(\mathbb{R}/I_{v,w}, t) = 1.$$

- *Otherwise, let i be a right ascent of v , so $vs_i > v$. Then*

- (1) *If i is a descent of w , so $ws_i < w$, then*

$$\mathcal{K}(\mathbb{R}/I_{v,w}, t) = \mathcal{K}(\mathbb{R}/I_{vs_i, w}, t).$$

- (2) *If i is an ascent of w , so $ws_i > w$, then*

$$\begin{aligned} \mathcal{K}(\mathbb{R}/I_{v,w}, t) &= \mathcal{K}(\mathbb{R}/I_{vs_i, w}, t) + (1 - t_{v(i)}/t_{v(i+1)})\mathcal{K}(\mathbb{R}/I_{vs_i, ws_i}, t) \\ &\quad - (1 - t_{v(i)}/t_{v(i+1)})\mathcal{K}(\mathbb{R}/I_{vs_i, w}, t). \end{aligned}$$

Our strategy to prove Theorem 6.2 is to show that the K-polynomials under the usual action for the Reisner–Stanley rings $\mathbb{R}/K_{v,w}$ associated to the pipe complexes $\Delta_{v,w}$ satisfy the same recursion. Our proof for this fact parallels that of [Knu08] for subword complexes; we could in fact refer to [Knu08, Corollary 2] by showing that the grading on $\mathbb{R}$ given by the usual action matches the grading given in the cited Corollary. However, that matching demands about as much analysis as the direct argument we give below.

If $v \not\leq w$, then $\Delta_{v,w}$ is the empty complex (since $w_0v \not\leq w_0w$, so no subwords of a reduced word for w_0v can be a reduced word for w_0w). Thus, in this case,

$$\mathcal{K}(\mathbb{R}/K_{v,w}, t) = 0.$$

If $v = w = w_0$, then $\Delta_{v,w}$ is the simplicial complex (on zero vertices) whose only face is the empty face. Therefore, in this case

$$\mathcal{K}(\mathbb{R}/K_{v,w}, t) = 1.$$

Otherwise, we rely on the vertex decomposition of $\Delta_{v,w}$ given by Theorem 6.4. First note that the homeomorphisms of Claims 6.9 and 6.10 are weight preserving by the following argument. Boxes in $D(v)$ which are not in column i or $i+1$ remain in the same place, and for $k \neq i, i+1$, the variable z_{jk} has weight

$$t_{v(k)} - t_{n-j+1} = t_{vs_i(k)} - t_{n-j+1}$$

in both $\mathbb{C}[z^{(v)}]$ and $\mathbb{C}[z^{(vs_i)}]$. A box in the column i and some row j of $D(v)$ corresponds to the box in row j and column $i+1$ in $D(vs_i)$ (unless $n-j+1 = v(i+1)$, in which case the box is deleted). The weight of $z_{j,i}$ in $\mathbb{C}[z^{(v)}]$ is

$$t_{v(i)} - t_{n-j+1},$$

which is equal to

$$t_{vs_i(i+1)} - t_{n-j+1},$$

the weight of $z_{j,i+1}$ in $\mathbb{C}[z^{(vs_i)}]$.

Furthermore, the variable z_{last} corresponding to V is $z_{n-v(i+1)+1,i}$. Therefore, z_{last} has weight $t_{v(i)} - t_{v(i+1)}$.

A face of $\Delta_{v,w}$ either contains V or not. Therefore, it is either a face of $\text{del}_V(\Delta_{v,w})$, or the union of a face of $\text{link}_V(\Delta_{v,w})$ with V . Let $\rho(a)$ denote the weight of the variable associated to a vertex a . Now [MilStu05, Theorem 1.13] (with the appropriate substitution to account for our use of the usual action rather than the rescaling action) states that

$$\mathcal{K}(R/K_{v,w}, \mathbf{t}) = \sum_{\sigma \in \Delta_{v,w}} \left(\prod_{a \in \sigma} t^{\rho(a)} \cdot \prod_{a \notin \sigma} (1 - t^{\rho(a)}) \right).$$

When i is a descent of w , Theorem 5.4 (A) asserts

$$\text{link}_V(\Delta_{v,w}) = \text{del}_V(\Delta_{v,w}) = \Delta_{vs_i,w}.$$

Therefore

$$\begin{aligned} \mathcal{K}(R/K_{v,w}, \mathbf{t}) &= t_{v(i)}/t_{v(i+1)} \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}) + (1 - t_{v(i)}/t_{v(i+1)}) \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}) \\ &= \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}). \end{aligned}$$

When i is an ascent of w , Theorem 5.4 (B) states that

$$\text{link}_V(\Delta_{v,w}) = \Delta_{vs_i,w} \quad \text{and} \quad \text{del}_V(\Delta_{v,w}) = \Delta_{vs_i,ws_i}.$$

Therefore,

$$\begin{aligned} \mathcal{K}(R/K_{v,w}, \mathbf{t}) &= t_{v(i)}/t_{v(i+1)} \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}) + (1 - t_{v(i)}/t_{v(i+1)}) \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}) \\ &= \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}) + (1 - t_{v(i)}/t_{v(i+1)}) \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}) \\ &\quad - (1 - t_{v(i)}/t_{v(i+1)}) \mathcal{K}(R/K_{vs_i,w}, \mathbf{t}). \end{aligned}$$

Therefore, the K -polynomials for $R/K_{v,w}$ satisfy the same recurrence relations as the K -polynomials for $R/I_{v,w}$, and hence they are equal. $\square$

6.3. Proof of Theorem 6.1. The following is immediate from Lemma 6.5:

Definition–Lemma 6.13. Given $f \in \mathbb{C}[z^{(v_{s_i})}]$ let f° be obtained by the substitution $z_{j,i+1} \mapsto z_{j,i}$. Then $f^\circ \in \mathbb{C}[z^{(v)}]$.

In what follows, let $\prec$ refer to the lexicographic term order we use on $Z^{(v_{s_i})}$ and $\prec_\circ$ be the term order on $Z^{(v)}$; see Section 2.3 for a definition of this term order.

The following is clear:

Lemma 6.14. If $\mathcal{L}$ is the leading term of $f \in \mathbb{C}[z^{(v_{s_i})}]$ with respect to $\prec$, then $\mathcal{L}^\circ$ is the leading term of $f^\circ \in \mathbb{C}[z^{(v)}]$ with respect to $\prec_\circ$.

The main technical point of this paper is below:

Proposition 6.15. Let i be the last ascent of $v \in S_n$. Suppose

- (I) i is a descent of $w \in S_n$ and $\mathcal{L}$ is a leading term of an essential minor of $I_{v_{s_i}, w}$; or
- (II) i is an ascent of $w \in S_n$ and $\mathcal{L}$ is the leading term of an essential minor of $I_{v_{s_i}, w}$; or
- (III) i is an ascent of $w \in S_n$ and $\mathcal{L}$ is the leading term of an essential minor of $I_{v_{s_i}, w_{s_i}}$.

Then in cases (I) and (III), $\mathcal{L}^\circ$ is divisible by the leading term $\mathcal{L}'$ with respect to $\prec_\circ$ of an essential minor of $I_{v, w}$. In case (II), the same holds for $\mathcal{L}^\circ z_{\text{last}}$.

Proof of Proposition 6.15: The basic idea of the proof is as follows: given an essential determinant D in $I_{v_{s_i}, w}$ (or $I_{v_{s_i}, w_{s_i}}$) which uses the submatrix M of $Z^{(v_{s_i})}$ and has leading term $\mathcal{L}$, we consider a determinant D' of $I_{v, w}$ that uses the submatrix M' of $Z^{(v)}$ with the same columns and rows as M except that if column i is used in M , we use column $i + 1$ in M' , and vice versa. In view of Lemma 6.5, *usually* this works to give an essential determinant

$$D' = \det M'$$

whose leading term $\mathcal{L}'$ has the desired properties. However, this sometimes fails, and our analysis below accounts for this. Each of (I), (II) and (III) is handled in four subcases, depending on which of the i -th and $i + 1$ -th columns of $Z^{(v_{s_i})}$ M uses.

Case I.1 (M uses neither the i -th nor $i + 1$ -th column): Let M' be the submatrix that uses the same rows and columns as M . By Lemma 6.5, $Z^{(v_{s_i})}$ and $Z^{(v)}$ do not differ in these columns, so we have that

$$D^\circ = D = D'.$$

Hence by Lemma 6.14,

$$\mathcal{L}^\circ = \mathcal{L} = \mathcal{L}',$$

so in particular $\mathcal{L}'$ divides $\mathcal{L}^\circ$.

Case I.2 (M uses both the i -th and $i + 1$ -th column): We may assume $D \neq 0$, hence:

Claim 6.16. The only nonzero entry of M in the i -th column comes from row $v_{s_i}(i) = v(i + 1)$ of $Z^{(v_{s_i})}$, and that entry is a 1.

Proof. Follows from Lemma 6.5 and the hypothesis that i is the last ascent of v . □

First construct a submatrix M'' of $Z^{(vs_i)}$ by changing M by replacing the 0 at position $(vs_i(i), i + 1)$ by $z_{n-v(i+1)+1, i+1}$ and switching the i -th and $i + 1$ -th columns. Now, we can compute D by cofactor expansion along the i -th column, so, by Claim 6.16, changing any entry in row $vs_i(i)$ of M other than the 1 in column i does not change the determinant. Hence

$$D'' = \pm D.$$

Let M' be the submatrix of $Z^{(v)}$ that uses the same rows and columns as M . By Lemma 6.5,

$$(D'')^\circ = \pm D^\circ$$

equals the essential determinant D' . Thus by Lemma 6.14,

$$\mathcal{L}' = \pm \mathcal{L}^\circ,$$

and therefore $\mathcal{L}'$ divides $\mathcal{L}^\circ$.

Case I.3 (M uses the i -th column but not the $i + 1$ -th column): M appears as a submatrix M' (with determinant D' and leading term $\mathcal{L}'$) of $Z^{(v)}$ using the same rows and columns except that the i -th column is replaced by the $i + 1$ -th. Therefore, by Lemma 6.5,

$$D = D' = D^\circ$$

and

$$\mathcal{L} = \mathcal{L}' = \mathcal{L}^\circ,$$

so $\mathcal{L}'$ divides $\mathcal{L}^\circ$. Thus we are done in this case provided D' is an essential minor of $I_{v,w}$, which follows from:

Claim 6.17. *If i is a descent of w then there are no boxes of the essential set of $D(w)$ in column i .*

Proof. Any box of $D(w)$ in column i must have a box of $D(w)$ to its immediate right. $\square$

Case I.4 (M uses the $i + 1$ -th column but not the i -th column): If M does not use row $vs_i(i) = v(i + 1)$ then let M' be the submatrix of $Z^{(v)}$ that uses the same rows and columns of M except that we use column i instead of column $i + 1$. Then by Lemma 6.5 it follows

$$D^\circ = D'.$$

Hence, by Lemma 6.14,

$$\mathcal{L}^\circ = \mathcal{L}'.$$

Moreover, D' is still essential since M' uses columns weakly to the left of those of M (and uses the same rows).

On the other hand, if M uses row $vs_i(i) = v(i + 1)$, let j be the column of the entry in the leading term of D used from row $v(i + 1)$. Construct a submatrix M'' of $Z^{(vs_i)}$ by replacing column j with column i (with associated determinant D'' and leading term $\mathcal{L}''$ with respect to $\prec$). Note that column i is to the right of column j and consists only of 0's except for a 1 in row $v(i + 1)$. Now, since by assumption M uses column $i + 1$, D'' is essential for w . It is easy to see from the fact $\prec$ is a lexicographic order that

$$\mathcal{L}'' = \pm \mathcal{L} / z_{n-v(i+1)+1, j}.$$

Now let M' be the submatrix of $Z^{(v)}$ using the same rows and columns as M'' . Then noting that M'' uses both columns i and $i + 1$, we can repeat the argument of Case I.2 to see that

$$\mathcal{L}' = \pm (\mathcal{L}'')^\circ = \pm (\mathcal{L} / z_{n-v(i+1)+1, j})^\circ = \pm \mathcal{L}^\circ / z_{n-v(i+1)+1, j}.$$

Hence, $\mathcal{L}'$ divides $\mathcal{L}^\circ$ as desired.

Case II.1 (M uses neither the i -th nor $i + 1$ -th column): This is proved exactly as in Case I.1.

Case II.2 (M uses both the i -th and $i + 1$ -th columns): This is proved exactly as in Case I.2.

Case II.3 (M uses only the i -th column but not the $i + 1$ -th column): Construct a submatrix M' of $Z^{(v)}$ by taking M and replacing the i -th column of $Z^{(v_{si})}$ with the i -th column of $Z^{(v)}$ and leaving all other columns of M unchanged. Note that

$$\mathcal{L}^\circ = \mathcal{L}.$$

However, $\mathcal{L}'$ may not divide $\mathcal{L}^\circ$. Instead we wish to prove

$$\mathcal{L}' = \pm \mathcal{L}^\circ z_{\text{last}}$$

(and hence $\mathcal{L}'$ divides $\mathcal{L}^\circ z_{\text{last}}$).

We assert that the position in each column of M' that contributes to $\mathcal{L}'$ is the same as for $\mathcal{L}$ (except that we use z_{last} in column i rather than 1, respectively). This is straightforward. Clearly D' is essential for $I_{v,w}$ since it uses the same rows and columns as D .

Case II.4 (M uses the $i + 1$ -th column and not the i -th column): This is proved exactly as in Case I.4.

In the analysis of (III), the main new issue is that we must show that, given an essential minor D for ws_i , the newly constructed minor D' is essential for w instead.

Case III.1 (M uses neither the i -th nor $i + 1$ -th column): The argument given in Case I.1 constructs a determinant D' such that $\mathcal{L}'$ divides $\mathcal{L}^\circ$. It remains to show that D' is essential. The only places where the rank matrices R^{ws_i} and R^w differ are in column i and rows t for which

$$ws_i(i + 1) < t \leq ws_i(i).$$

Note moreover that no boxes of $D(ws_i)$ lie in this region. Now, let d be the essential set box of ws_i causing D to be essential. If d is not in column $i + 1$, or d is in column $i + 1$ and strictly south of row $ws_i(i)$, it easily follows that D' is also essential for w . Otherwise d must be in position

$$(ws_i(i + 1) + 1, i + 1).$$

Then d is no longer even a box of $D(w)$, but the box d^* in position

$$(ws_i(i + 1) + 1, i)$$

(to the immediate left of d) is in $\mathcal{E}(w)$. In addition,

$$R_{d^*}^w = R_d^{ws_i} - 1,$$

and the columns of D are weakly to the left of column i .

Consider the positions of the variables in M contributing to the leading term $\mathcal{L}$ of D . Let M'' be the submatrix defined by any $R_{d^*}^w = R_d^{ws_i} - 1$ of these positions. (For definiteness, we can take all but the rightmost position, that of variable $z_{\text{rightmost}}$.) Then M'' has leading term $\mathcal{L}''$ given by the product of the aforementioned variables we picked out, so

$$\mathcal{L} = \mathcal{L}'' z_{\text{rightmost}}.$$

(Otherwise we would contradict the fact that $\mathcal{L}$ is a leading term of D .) Now let M' be the submatrix of $Z^{(vs_i)}$ using the same rows and columns as M'' . Then

$$\mathcal{L}' = \mathcal{L}'' = (\mathcal{L}'')^\circ,$$

and it follows that

$$\mathcal{L}' = \mathcal{L}^\circ z_{\text{rightmost}},$$

and hence $\mathcal{L}'$ divides $\mathcal{L}^\circ$.

Case III.2 (M uses both the i -th and $i + 1$ -th column): Repeat the construction of Case I.2 to obtain a minor D'' . Note that since M uses column $i + 1$, the essential set box d of ws_i causing D to be essential is weakly to the right of column $i + 1$. If d is strictly to the right, then let D' be the determinant of $Z^{(v)}$ that uses the same rows and columns as M . This case follows as in Case I.2. Otherwise, if d lies in column $i + 1$ and is at

$$(ws_i(i + 1) + 1, i + 1),$$

the box d^* to its immediate left is an essential box for w , of rank one less, as in Case III.1. Now let M' be the one smaller minor that uses the same rows and columns as M , except that it excludes row $v(i + 1)$ and column $i + 1$. Then D' is essential for w due to box d^* . This case then follows.

Case III.3 (M uses column i but not column $i + 1$): To construct D' we use the same construction as in Case I.3, and its essentialness follows as in Case III.2.

Case III.4 (M uses column $i + 1$ but not column i): Use the same construction as in Case I.4. There are two cases to prove essentialness, paralleling the two subcases of Case I.4. If M does not use row $vs_i(i)$ then we apply the essential box argument of Case III.1. In the other subcase, we argue essentialness as in Case III.2. $\square$

Conclusion of the proof of Theorem 6.1: We induct on $\ell(w_0v) \geq 0$. The base case of $\ell(w_0v) = 0$, which is where $v = w_0$, is trivial since $z^{(v)} = \emptyset$. For the induction step, assume that $\ell(w_0v) \geq 1$; hence $z^{(v)} \neq \emptyset$, and in particular, we have a last variable z_{last} and associated vertex $V \in \Delta_{v,w}$, as defined above.

Let $\mathcal{P}$ be a nonface of $\Delta_{v,w}$; we must show $z^{\mathcal{P}}$ is divisible by the leading term of a defining minor of $I_{v,w}$.

Suppose $V \in \mathcal{P}$. Then (1) of Lemma 6.3 asserts $\mathcal{P} \setminus V$ is a nonface of $\text{link}_V(\Delta)$. There are then two cases, depending on whether i is a descent or ascent of w . If i is a descent of w , then part (A) of Theorem 6.4 says

$$\text{link}_V(\Delta) \cong \Delta_{vs_i,w}.$$

Under the relabeling map of Claim 6.9, $\tilde{\mathcal{P}} \setminus V$ is a nonface of $\Delta_{vs_i,w}$ and hence by induction $z^{\tilde{\mathcal{P}} \setminus V}$ is divisible by the leading term of a defining minor of $I_{vs_i,w}$. The conclusion then follows from part (I) of Proposition 6.15. Similar arguments apply for the case where $V \in \mathcal{P}$ and i is an ascent of w , and the two cases (depending on whether i is a descent or ascent of w) where $V \notin \mathcal{P}$. $\square$

ACKNOWLEDGEMENTS

We thank Allen Knutson for a number of inspiring communications and suggestions and in particular for sharing his observation about homogeneity and multiplicities of Schubert varieties. We also thank Rebecca Goldin, Li Li, Hal Schenck and Frank Sottile for helpful conversations. This paper was partially completed during the NSF VIGRE supported meeting on “Combinatorial Algebraic Geometry of Flag Varieties” at the University of Iowa; we thank the organizers Megumi Harada and Julianna Tymoczko. AY is partially supported by NSF grants DMS-0601010 and DMS-0901331. We made extensive use of the computer algebra system Macaulay 2 during our investigations.

REFERENCES

- [BerBil93] N. Bergeron and S. Billey, *RC-graphs and Schubert polynomials*, Experiment. Math. **2**(1993), no. 4, 257–269.
- [BilPro79] L. J. Billera and J. S. Provan, *A decomposition property for simplicial complexes and its relation to diameters and shellings*, Second International Conference on Combinatorial Mathematics (New York, 1978), New York Acad. Sci., New York, 1979, pp. 82–85.
- [Bil99] S. Billey, *Kostant polynomials and the cohomology ring for G/B* , Duke Math. J. **96**(1999), no. 1, 205–224.
- [BruHer98] W. Bruns and J. Herzog, *Cohen–Macaulay rings, Revised ed.*, Cambridge studies in advanced mathematics **39**. Cambridge University Press, Cambridge, 1998.
- [BruVet88] W. Bruns and U. Vetter, *Determinantal rings*, Monografías de Matemática, **45**. Instituto de Matemática Pura e Aplicada, Rio de Janeiro, 1998.
- [BucRim04] A. Buch and R. Rimányi, *Specializations of Grothendieck polynomials*, C. R. Acad. Sci. Paris, Ser. I **339**(2004), 1–4.
- [FomKir94] S. Fomin and A. N. Kirillov, *Grothendieck polynomials and the Yang-Baxter equation*, Proc. 6th Intern. Conf. on Formal Power Series and Algebraic Combinatorics, DIMACS, 1994, 183–190.
- [FomKir96] ———, *Combinatorial B_n -analogues of Schubert polynomials*, Trans. A. M. S., **348**(1996), 3591–3620.
- [Ful92] W. Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), no. 3, 381–420.
- [Ful97] ———, *Young tableaux*, London Mathematical Society Student Texts, 35. Cambridge University Press, Cambridge, 1997.
- [GhoRag06] S. Ghorpade and K. N. Raghavan, *Hilbert functions of points on Schubert varieties in the symplectic Grassmannian*, Trans. Amer. Math. Soc. **358** (2006), 5401–5423.
- [Gol01] R. F. Goldin, *The cohomology ring of weight varieties and polygon spaces*, Adv. Math. **160**(2001), no. 2, 175–204.
- [HamPit08] A. Hammett and B. Pittel, *How often are two permutations comparable?* Trans. A. M. S. **360**(2008), no. 9, 4541–4568.
- [Hoc77] M. Hochster, *Cohen-Macaulay Rings, combinatorics, and simplicial complexes*, Ring theory, II (Proc. Second Conf., Univ. Oklahoma, Norman, OK, 1975) (B.R. Macdonald and R. Morris, eds.) Lecture Notes in Pure and Applied Mathematics **26**, Marcel Dekker, New York (1977), 171–223.
- [IkeNar07] T. Ikeda and H. Naruse, *Excited Young diagrams and equivariant Schubert calculus*, preprint 2009. arXiv:math.AG/07030637
- [KazLus79] D. Kazhdan and G. Lusztig, *Representations of Coxeter Groups and Hecke Algebras*, Invent. Math. **53** (1979), 165–184.
- [Knu07] A. Knutson, *Automatically reduced degenerations of automatically normal varieties*, preprint, 2007. arXiv:0709.3999
- [Knu08] ———, *Schubert patches degenerate to subword complexes*, Transform. Groups **13** (2008), 715–726.
- [Knu09] ———, *Frobenius splitting and degeneration, especially of Schubert varieties*, preprint, 2009.

- [KnuMil04] A. Knutson and E. Miller, *Subword complexes in Coxeter groups*, Adv. Math. **184**(2004), no. 1, 161–176.
- [KnuMil05] ———, *Gröbner geometry of Schubert polynomials*, Ann. of Math. (2) **161** (2005), no. 3, 1245–1318.
- [KnuMilYon05] A. Knutson, E. Miller and A. Yong, *Gröbner geometry of vertex decompositions, and of flagged tableaux*, J. Reine Agnew Math., to appear.
- [KosKum90] B. Kostant and S. Kumar, *T-equivariant K-theory of generalized flag varieties*, J. Differential Geom. **32**(1990), no. 2, 549–603.
- [Kra01] C. Krattenthaler, *On multiplicities of points on Schubert varieties in Grassmannians*, Séminaire Lotharingien Combin. **45**(2001), Article B45c, 11 pp.
- [Kre08] V. Kreiman, *Schubert classes in the equivariant K-theory and equivariant cohomology of the Grassmannian*, preprint 2008.
- [KreLak04] V. Kreiman and V. Lakshmibai, *Multiplicities at Singular Points of Schubert Varieties in the Grassmannian*, Algebra, arithmetic and geometry with applications, 553–563, Springer, Berlin, 2004.
- [LakWey90] V. Lakshmibai and J. Weyman, *Multiplicities of points on a Schubert variety in a minuscule G/P* , Adv. Math. **84**(1990), no. 2, 179–208.
- [LasSch82a] A. Lascoux and M. P. Schützenberger, *Polynômes de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), no. 13, 447–450.
- [LasSch82b] ———, *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 11m 629–633.
- [Man01] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*, American Mathematical Society, Providence 2001.
- [MilStu05] E. Miller and B. Sturmfels, *Combinatorial commutative algebra*, Springer-Verlag, 2005.
- [Mir07] R. Miró-Roig, *Determinantal ideals*, Progress in Mathematics, **264**, Birkhäuser Verlag, Basel, 2008.
- [RagUpa07] K. N. Raghavan and S. Upadhyay, *Hilbert functions of points on Schubert varieties in the orthogonal Grassmannian*, preprint, 2007. arXiv:0704.0542
- [RosZel01] J. Rosenthal and A. Zelevinsky, *Multiplicities of points on Schubert varieties in Grassmannians*, J. Algebraic Combinatorics, **13**(2001), 213–218.
- [Ste01] J. Stembridge, *Minuscule elements of Weyl groups*, J. Algebra **235** (2001), 722–743.
- [Wil06] M. Willems, *K-théorie équivariante des tours de Bott. Application à la structure multiplicative de la K-théorie équivariante des variétés de drapeaux*, Duke Math. J. **132**(2006), no. 2, 271–309.
- [WooYon08] A. Woo and A. Yong, *Governing singularities of Schubert varieties*, J. Algebra, **320**(2008), no. 2, 495–520.

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, SAINT OLAF COLLEGE, NORTH-FIELD, MN 55057

E-mail address: woo@stolaf.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801

E-mail address: ayong@illinois.edu